


PROVINCE OF THE EASTERN CAPE
IPHONDO LEMPUMA KOLONI
PROVINSIE OOS-KAAP

# Provincial Gazette Igazethi Yephondo Provinsiale Koerant

Vol. 21 KING WILLIAM'S TOWN, 3 JUNE 2014

No. 3191 (Extraordinary)

## We all have the power to prevent AIDS



Prevention is the cure

AIDS HEWUNE

0800 012 322

DEPARTMENT OF HEALTH

N.B. The Government Printing Works will not be held responsible for the quality of "Hard Copies" or "Electronic Files" submitted for publication purposes





9771682455006

#### **IMPORTANT NOTICE**

The Government Printing Works will not be held responsible for faxed documents not received due to errors on the fax machine or faxes received which are unclear or incomplete. Please be advised that an "OK" slip, received from a fax machine, will not be accepted as proof that documents were received by the GPW for printing. If documents are faxed to the GPW it will be the sender's responsibility to phone and confirm that the documents were received in good order.

Furthermore the Government Printing Works will also not be held responsible for cancellations and amendments which have not been done on original documents received from clients.

#### **CONTENTS • INHOUD**

No. Page Gazette

#### PROVINCIAL NOTICE

#### PROVINCIAL NOTICE

No. 30

# PROVINCE OF THE EASTERN CAPE PROVINCIAL NOTICE DEPARTMENT OF ECONOMIC DEVELOPMENT AND ENVIRONMENTAL AFFAIRS

PUBLISHING OF THE NELSON MANDELA BAY MUNICIPALITY AS A BIOREGION IN TERMS OF SECTION 40 (1) OF THE NATIONAL ENVIRONMENTAL MANAGEMENT: BIODIVERSITY ACT, 2004 (ACT NO 10 OF 2004) AND INVITATION TO COMMENT ON THE DRAFT BIOREGIONAL PLAN FOR THE NELSON MANDELA BAY MUNICIPALITY AS PER SECTIONS 47 (2), 99 (1), 99 (2)(C), 100 (1)(a) AND 100(2).

I, MCEBISI JONAS, Member of the Executive Council responsible for Economic Development, Environmental Affairs and Tourism in the Province of the Eastern Cape, acting in terms of section 40 (1) gives notice to publish in the Gazette, Nelson Mandela Bay Municipality as a Bioregion in terms of Section 40 (1) (a) (b). I hereby also invite comment on the Draft Bioregional Plan for the Nelson Mandela Bay Municipality as per Sections 47 (2), 99 (1), 99 (2)(c), 100 (1)(a) and 100 (2). of the National Environmental Management: Biodiversity Act, 2004 (Act No 10 of 2004)

The boundary of the bioregion for the bioregional plan is the boundary of the Nelson Mandela Bay Municipality (NMBM) in the Eastern Cape Province of South Africa.

Copy of Bioregional Plan can be viewed on the Nelson Mandela Bay Municipality website <a href="https://www.nelsonmandelabay.gov.za">www.nelsonmandelabay.gov.za</a>

All comments and enquiries must be forwarded to Mr Albert Mfenyana within 30 days of publication of this notice via email <u>Albert.Mfenyana@deaet.ecape.gov.za</u>, fax 043 605 7300. or

**By post to:**General Manager
Environmental Affairs

Private Bag X0054

Bisho 5605 By hand at:

2<sup>nd</sup> floor, Room 288

Beacon Hill

Corner of Hargreaves Street & Hockley Close

King Williams Town

5600

M. JONAS

MEC FOR ECONOMIC DEVELOPMENT, ENVIRONMENTAL AFFAIRS AND TOURISM

# Nelson Mandela Bay Municipality Draft Bioregional Plan

Nelson Mandela Bay Municipality
Public Health Directorate
Environmental Management Sub-Directorate

Report No 400919/2

Page i

#### **Executive Summary**

This draft bioregional plan relates to the Nelson Mandela Bay Municipality in the Eastern Cape Province of South Africa. As this is the first bioregional plan for the Nelson Mandela Bay Municipality it does not replace any previously published Bioregional Plan.

The purpose of the draft bioregional plan is to provide a map of biodiversity priorities and accompanying guidelines to inform land-use planning, environmental assessment and authorisations, and natural resource management by a range of sectors whose policies and decisions impact on biodiversity.

The bioregional plan is a spatial plan that shows terrestrial and aquatic features that are critical for conserving biodiversity and maintaining ecosystem functioning. These areas are referred to as Critical Biodiversity Areas and Ecological Support Areas. In addition, the bioregional plan provides a profile of priority biodiversity in the bioregion, outlines other measures for effective management of biodiversity, and includes recommendations for review, monitoring and updating. The bioregional plan is underpinned by the *Conservation Assessment and Plan for the Nelson Mandela Bay Municipality* (SRK Consulting, 2009), a systematic biodiversity plan that was developed according to established protocols.

It is important to note that the conservation assessment was developed in a manner that attempts to minimise potential conflict between biodiversity and other forms of land-use to the greatest extent possible. Consequently, the draft bioregional plan is consistent with the National Environmental Management: Biodiversity Act (Act No. 10 of 2004), and meets all the requirements of the *Guideline regarding the Determination of Bioregions and the Preparation and Publication of Bioregional Plans* (DEAT, 2008) for review by a Ministerial Panel.

The National Environmental Management Act (NEMA; Act No. 107 of 1998) requires that all organs of state must take biodiversity considerations into account in their decision-making. The bioregional plan facilitates this by serving as the official reference for biodiversity priorities within the bioregion, superseding all the previous non-published spatial biodiversity plans.

The plan supports the principles of integrated development planning and sustainable development espoused by the NEMA by identifying the biodiversity conservation priority in the municipal area, as well as other natural areas that could potentially be made available for other land-uses. The plan is consistent with the NEM: Biodiversity Act (Act No. 10 of 2004), the National Biodiversity Framework (2009), and South Africa's commitments in terms of international agreements such as the Convention on Biological Diversity (1993) and the United Nations Framework Convention on Climate Change. Furthermore, the plan was developed in an integrated manner with the Municipal Spatial Development Framework (SDF), which is a spatial articulation of the Municipal Integrated Development Plan (IDP), in order to identify an optimal balance between biodiversity conservation and other sector needs.

The objectives of the bioregional plan are to:

- Form the primary biodiversity informant for land-use and resource use decision-making;
- Identify a network of Critical Biodiversity Areas that achieve national biodiversity thresholds on the least amount of land possible; and

STEW/reet

Page ii

Act as the spatial framework and policy for sustainable development and to assist the municipality
in complying with environmental and planning legislation requirements which promote the
protection and management of biodiversity.

The bioregional plan should be used by all sectors that are involved in land-use planning and decision-making and multi-sectoral planning. The users of the bioregional plan include mandatory users, i.e. those who are compelled to consider the bioregional plan, and other intended users for whom the bioregional plan will be a useful planning and developmental tool. The bioregional plan has three main uses:

- Reactive decision-making, such as environmental impact assessment, agricultural land-use decisions and development control decisions through land-use legislation (e.g. rezoning, planning approvals etc.);
- 2. Proactive forward planning, such as IDPs, SDFs and zoning schemes; and
- 3. Proactive conservation, such as biodiversity stewardship initiatives and the expansion of protected areas.

Page iii

#### **Acknowledgements**

The authors would like to acknowledge the following contributors to, or reviews of, the Bioregional Plan and Systematic Conservation Assessment and Plan for the Nelson Mandela Bay Municipality:

Joram Mkosana, Abigail Kamineth, Stuart Beattie, Schalk Potgieter, Jill Miller, Nadia Wessels, Alan de Vries, Wesley Berrington, Jeff Manuel, Dr Tammy Smith, Dr. Stephen Holness, Wilma Lutsch, Kiruben Naicker, Rob Gardiner, Prof. Richard Cowling, Dr Mandy Lombard, Dr Paul Martin, Dr Dave Balfour, Dr Derek du Preez, Dawn McCarthy, Kithi Ngesi, Adri Scholtz, the NMBM Drawing Office staff, Albert Mfenyana, Phumla Mzazi-Geja, Noluthando Bam, Lulama Macanda, Andries Struwig, Nicholas Scarr, Alan Southwood, the Table Mountain Fund, Dr Matthieu Rouget, Deryk Langman, Denise Davies, Arthur Rump, Eckart Schumann, An van Cauter, Rebecca Sims-Castley, Dr Bill Branch, Lloyd Wingate, Fred Kigozi, Jamie Pote, Kristal Maze, Mark Botha, Amrei von Hase, Andrew Knight, Tania Katzschner, Dr Mandy Cadman, Dr Debra Roberts and Richard Boon.

Page iv

#### **Table of Contents**

	⊨xe	cutive	Summary	
	Ack	nowled	gements	ii
1	Pari	t A: Inf	roduction and Objectives	g
	1.1	Purpo	ose and Objectives of the Bioregional Plan	g
	1.2	Inten	ded Users and Uses of the Bioregional Plan	10
	1.3	Limita	ations	12
2	Part	t B: Bi	odiversity Profile	13
	2.1	Signit	icance of the Biodiversity within the Bioregion	13
	2.2	Impo	tant Terrestrial Features	14
	2.3	Impo	tant Freshwater Features	18
	2.4	Impo	rtant Coastal Features and Estuaries	18
	2.5	Impo	tant Ecological Corridors	19
	2.6	Patte	rns of Land-Use	19
	2.7	Ecos	ystem Protection Level within the Nelson Mandela Bay Municipal Area	21
	2.8	Desc	ription of Critical Biodiversity Areas	21
3	Pari	t C: Cr	itical Biodiversity Areas map	24
4	Part	t D: Gu	ıidelines	25
4 5			her measures for effective management	
		t E: Ot		43
	Part	t E: Ot	her measures for effective management	<b>43</b> 43
	Part	t <b>E: Ot</b> Plann	her measures for effective management	43 43
	Part	E: Ot Planr 5.1.1 5.1.2	her measures for effective managementing	43 43 45
	<b>Par</b> <sup>1</sup> 5.1	E: Ot Planr 5.1.1 5.1.2	her measures for effective management  ing  Protected Area Expansion  Ecosystems and Species Requiring Biodiversity Management Plans  ation and Management	434345
	<b>Par</b> <sup>1</sup> 5.1	Plann 5.1.1 5.1.2 Opera 5.2.1	her measures for effective management  ing	43434546
	<b>Par</b> <sup>1</sup> 5.1	Plann 5.1.1 5.1.2 Opera 5.2.1 5.2.2	her measures for effective management	4343454646
	<b>Part</b> 5.1 5.2	Plann 5.1.1 5.1.2 Opera 5.2.1 5.2.2	her measures for effective management  ing  Protected Area Expansion  Ecosystems and Species Requiring Biodiversity Management Plans  ation and Management  Invasive Alien Species Management  Priority Areas for Rehabilitation and Restoration	4345464647
	<b>Part</b> 5.1 5.2	Plann 5.1.1 5.1.2 Opera 5.2.1 5.2.2 Mains 5.3.1	her measures for effective management  ing  Protected Area Expansion  Ecosystems and Species Requiring Biodiversity Management Plans  ation and Management  Invasive Alien Species Management  Priority Areas for Rehabilitation and Restoration	4345464647
	<b>Part</b> 5.1 5.2 5.3	Plann 5.1.1 5.1.2 Opera 5.2.1 5.2.2 Mains 5.3.1	her measures for effective management  ling	434546474748
	<b>Part</b> 5.1 5.2 5.3	Plann 5.1.1 5.1.2 Opera 5.2.1 5.2.2 Mains 5.3.1 Buildi 5.4.1 5.4.2	her measures for effective management	43454647474848
	<b>Part</b> 5.1 5.2 5.3	Plann 5.1.1 5.1.2 Opera 5.2.1 5.2.2 Mains 5.3.1 Buildi 5.4.1 5.4.2 5.4.3	her measures for effective management	43454647474848
	<b>Part</b> 5.1 5.2 5.3	Plann 5.1.1 5.1.2 Opera 5.2.1 5.2.2 Mains 5.3.1 Buildi 5.4.1 5.4.2	her measures for effective management	4345464747484849

STEW/reeb

Draft Bioregional Plan 2 Dec 10 for gazetting (2014-final)

		sulting egional	Plan F	age v
		5.4.6	Payment for Ecosystem Services	 50
		5.4.7	Expanded Public Works Programme	50
6	Part	F: Mo	onitoring, Reviewing and Updating	51
	6.1	Revie	ew	51
	6.2	Monit	toring	51
		6.2.1	Monitoring Instruments	52
		6.2.2	Monitoring Indicators	52
	6.3	Upda	ating	53
7	Part	G: GI	IS Files	53
8	Con	clusic	ons and Recommendations	54
9	Refe	erence	es	56
Аp	pend	lices .		58
	Арр		A: Systematic Conservation Assessment and Plan for the Nelson Mandela cipality (SRK Consulting, 2010)	•
	Арр	endix l	B: Extensive Biodiversity Profile	59
	Арр	endix (	C: A3 map of Critical Biodiversity Areas	60
Li	st c	of Ta	ables	
Та	ble 1	-1: The	e mandatory and recommended users of the bioregional plan	10
Та	ble 2	-1: Key	y to Figure 2-3 – vegetation type numbers	15
Та	ble 2		overview of the Ecosystem Threat Status of the vegetation types of the NI	
Та	ble 2	-3: Dis	stribution of conservation areas within the municipal area	21
Та	ble 2	-4: Des	scription of the Critical Biodiversity Area categories	22
Та	ble 4		Biodiversity-compatible land-use guidelines matrix - Category descriptions anagement objectives	
Та	ble 4		Recommended land & resource (water) use management guidelines in Crodiversity Areas and other categories on the CBA map	
Та	ble 4	-3: Def	finitions of biodiversity-compatible land-use types	35
Та	ble 4	-4: Def	finitions of moderate impact rural land-use types	38
Та	ble 4	-5: Def	finitions of high impact agricultural land-uses	40
Та	ble 4	-6: Def	finitions of high impact urban and industrial land-uses	42
Та	ble 7	-1: Out	tline of the shapefiles included in the data package	53

SRK	Consulting	
Draft	Bioregional	Plan

Page vi

List of	Figures
---------	---------

Figure 2-2: T	he biomes within Nelson Mandela Bay municipal area (Stewart <i>et al.</i> , 2008) 14
=	The historical distribution of natural vegetation types in the Nelson Mandela Barunicipal area (Stewart <i>et al.</i> , 2008)
•	The Ecosystem Threat Status of vegetation types within the Nelson Mandela Bayunicipal area (Stewart <i>et al.</i> , 2008)
•	atterns of land-use in the Nelson Mandela Bay municipal area (SRK Consulting, 2009
· ·	nplementation sites in the Nelson Mandela Bay municipal area (SRK Consulting, 2009
Figure 3-1: M	ap of Critical Biodiversity Areas within the municipal area24
List of al	obreviations
C.A.P.E.	Cape Action for People and the Environment Programme
CBA	Critical Biodiversity Area
CBD	Convention on Biological Diversity
CDF	Conservation Development Framework
ESA	Ecological Support Area
CETT	Corporate EIA Task Team
DEA	Department of Environmental Affairs (National)
DEAT	former Department of Environmental Affairs and Tourism (National)
DEDEA	Department of Economic Development and Environmental Affairs (Eastern Cape Province)
DMR	Department of Mineral Resources
DWA	Department of Water Affairs
ECBCP	Eastern Cape Biodiversity Conservation Plan
EC Parks	Eastern Cape Parks
EAP	Environmental Assessment Practitioner
EIA	Environmental Impact Assessment
EIP	Environmental Implementation Plan
EMF	Environmental Management Framework
EMP	Environmental Management Plan or Programme

STEW/reeb

Draft Bioregional Plan 2 Dec 10 for gazetting (2014-final)

SRK Consulting

Draft Bioregional Plan Page vii

**ESA** Ecosystem Support Area

**EPWP** Extended Public Works Programme

IDP Integrated Development Plan

**MDTTT** Multi-Disciplinary Technical Task Team

**NEMA** National Environmental Management Act, 1998 (Act No. 107 of 1998)

**NMBM** Nelson Mandela Bay Municipality

**SANBI** South African National Biodiversity Institute

**SEA** Strategic Environmental Assessment

SDF Spatial Development Framework

**STEP** Subtropical Thicket Ecosystem Programme

#### Glossary

**Biodiversity** The variability among living organisms from all sources including, terrestrial, marine and

other aquatic ecosystems and the ecological complexes of which they are part and also

includes diversity within species, between species, and of ecosystems.

**Biome** A biome is a regional-scale ecosystem characterized by distinct types of vegetation, animals,

and microbes that have developed under specific soil and climatic conditions.

C.A.P.E. Cape Action for People and the Environment Programme is a multi-party government and

civil society initiative to conserve biodiversity, while creating benefits for all the people of

the Cape Floristic Region.

CoastCare The CoastCare programme consists of projects and products that contribute to the goals and

objectives of the coastal policy.

Ecosystem A dynamic system of plant, animal and micro-organism communities and their nonliving

environment interacting as a functioning unit.

Ecosystem The extent to which the biodiversity target for a vegetation type is conserved via protected

areas.

Protection

Level

The classification of vegetation types based on the extent of remaining area (currently not Ecosystem

transformed) of each vegetation type in relation to their biodiversity target.

Threat Status

LandCare LandCare is a community based and government supported approach to the sustainable

management and use of agricultural natural resources. The overall goal of LandCare is to

Page viii

optimise productivity and sustainability of natural resources so as to result in greater productivity, food security, job creation and better quality of life for all.

STEP

Subtropical Thicket Ecosystem Programme is a multi-party government and civil society initiative to conserve biodiversity, while creating benefits for all the people of the Subtropical Thicket biome.

Sustainable development

The integration of social, economic and environmental factors into planning, implementation and decision-making so as to ensure that development serves present and future generations.

Page 9

### 1 Part A: Introduction and Objectives

The boundary of the bioregion for this bioregional plan is the boundary of the Nelson Mandela Bay Municipality (NMBM) in the Eastern Cape Province of South Africa. Since this is the first bioregional plan for the NMBM, it does not replace any previously published bioregional plans.

#### 1.1 Purpose and Objectives of the Bioregional Plan

The purpose of the draft bioregional plan is to provide a map of biodiversity priorities and accompanying guidelines to inform land-use planning, environmental assessment and authorisations, and natural resource management by a range of sectors whose policies and decisions impact on biodiversity. The bioregional plan is a spatial plan that shows terrestrial and aquatic features that are critical for conserving biodiversity and maintaining ecosystem functioning. These areas are referred to as Critical Biodiversity Areas (CBAs).

The National Environmental Management Act (NEMA; Act No. 107 of 1998) requires that all organs of state take biodiversity considerations into account in their decision-making. The bioregional plan facilitates this by serving as the official reference for biodiversity priorities within the bioregion, superseding all the previous non-published spatial biodiversity plans<sup>1</sup>.

The plan acts as the spatial framework for sustainable development by identifying the minimum set of areas that are required for biodiversity conservation, the continuation of ecological processes and the provision of ecosystem goods and services. The plan thereby supports the principles of integrated development planning and sustainable development espoused by the NEMA. It is also consistent with the NEM: Biodiversity Act (Act No. 10 of 2004), the National Biodiversity Framework (2009), and South Africa's commitments in terms of international agreements such as the Convention on Biological Diversity (1993) and the United Nations Framework Convention on Climate Change.

The objectives of the bioregional plan are to:

- Form the primary biodiversity informant for land-use and resource use decision-making;
- Identify a network of Critical Biodiversity Areas that achieve national biodiversity thresholds on the least amount of land possible; and
- Act as the spatial framework and policy for sustainable development and to assist the municipality
  in complying with environmental and planning legislation requirements which promote the
  protection and management of biodiversity.

The bioregional is underpinned by the Conservation Assessment and Plan for the Nelson Mandela Bay Municipality (SRK Consulting, 2009). The NMBM conservation assessment is a systematic biodiversity plan that was developed according to established protocols. Systematic biodiversity

-

<sup>1</sup> The portions of the following plans pertaining to the NMBM are superseded by the NMBM Bioregional Plan: The Cape Action for People and the Environment (C.A.P.E.) (Cowling *et al.*, 1999 a & b); The Subtropical Thicket Ecosystem Plan (STEP) (Cowling *et al.*, 2003); and The Eastern Cape Biodiversity Conservation Plan (Berliner & Desmet, 2007).

Page 10

planning is a rigorous, data-driven approach for assessing the location, status and importance of a range of biodiversity features. It is widely accepted in South Africa, and internationally, as the best available science for identifying spatial biodiversity priority areas. It is important to note that the NMBM conservation assessment was developed in a way that attempts to minimise potential conflict between biodiversity and other forms of land-use.

#### 1.2 Intended Users and Uses of the Bioregional Plan

The bioregional plan should be used by all sectors that are involved in land-use planning and decision-making or multi-sectoral planning. The users of the bioregional plan include mandatory users, i.e. those who are compelled to consider the bioregional plan, and those who will find it to be a useful planning and developmental tool. The users of the bioregional plan are described in Table 1-1.

The bioregional plan has three main uses:

- 1. Reactive decision-making, such as environmental impact assessment and agricultural land-use decisions, water-use licensing, and development control decisions through land-use legislation (e.g. rezoning, planning approvals etc.);
- 2. Proactive forward planning, such as IDPs, SDFs and zoning schemes; and
- 3. Proactive conservation, such as biodiversity stewardship initiatives and the expansion of protected areas

Table 1-1: The mandatory and recommended users of the bioregional plan

User	Mandatory uses	Recommended uses
Nelson Mandela Bay Municipality	Must align the bioregional plan with its Integrated Development Plan (IDP) and Spatial Development Framework (SDF) and must demonstrate how the bioregional plan may be implemented. Must also take the bioregional plan into account in issuing planning authorisations.	Should integrate Critical Biodiversity Areas and other relevant guidelines and recommendations from the bioregional plan into Environmental Management Frameworks (EMFs) developed in terms of Chapter 8 of the EIA Regulations, and zoning schemes (also known as planning or land-use schemes).
Any organ of state that must prepare an Environmental Implementation Plan (EIP) or Environmental Management Plan (EMP) in terms of Chapter 3 of NEMA	Must consider the bioregional plan in developing the EIP or Environmental Management Plan.	-

Page 11

Uş	ser	Mandatory uses	Recommended uses
	Eastern Cape Department of Economic Development and Environmental Affairs (DEDEA)	Must consider the bioregional plan before issuing environmental authorisations.	Should take the bioregional plan into account when developing provincial supplements to the NEMA EIA regulations in terms of section 24A and in developing EMFs and Provincial Guidelines in terms of the Chapter 8 EIA regulations.  Should require that Environmental Assessment Practitioners (EAPs) refer to the bioregional plan in Basic Assessment Reports and Environmental Impact Reports.
Environmental	Eastern Cape Parks	-	Should use the bioregional plan to guide their inputs into land-use planning, land-use decision-making and natural resource management and to inform their management actions and the development of an Eastern Cape Protected Area Expansion Strategy.
decision-makers who are required by section 2(1)(c) of NEMA to apply the NEMA section 2	National Department of Agriculture, Forestry and Fisheries	Should take the bioregional plan into account in their authorisations (e.g. for ploughing virgin land and for the subdivision of agricultural land).	Should take the bioregional plan into account in their planning processes and in their programmes.
principles in their decision-making	National Department of Minerals and Energy	Should take the plan into account in their authorisations for prospecting and mining.	-
	Eastern Cape Department of Agriculture and Land Affairs	Should take the bioregional plan and identified Critical Biodiversity Areas into account in their comments on applications.	Should take bioregional plan into account in planning and scheduling LandCare activities, and in any other planning activities (such as farm planning and Area-Wide Planning)  Should take the bioregional plan into account in planning and implementing land reform programmes, and in the development of policy, legislation or guidelines for land-use planning and management.
	Eastern Cape Department of Public Works	-	Should take the bioregional plan into account when identifying locations and routes for roads and other infrastructure.

Page 12

December 2010

Us	ser	Mandatory uses	Recommended uses
	Eastern Cape Department of Housing, Local Government and Traditional Affairs	-	Should take the bioregional plan into account in development of the Provincial Growth and Development Strategy and when identifying suitable sites for new housing developments.
	Catchment Management Agencies	Should integrate the bioregional plan in their decision-making about water applications	-
Organs of state pre in terms of Reg. 74 regulations		-	Should take the bioregional plan into account
Working for Water, Working for Wetlands, LandCare, CoastCare, and other programmes that deal with maintaining and restoring natural resources		-	Should take the bioregional plan into account in planning and scheduling their activities
Environmental and planning consultants		-	Should take the bioregional plan into account when undertaking Strategic Environmental Assessments and Environmental Impact Assessments or when developing IDPs, SDFs or EMFs for a municipality
Conservation NGOs		-	Should use the bioregional plan to guide comments on planning tools such as SDFs and on development applications, and to direct on-the-ground conservation initiatives
Private landowners		-	Should use the bioregional plan if they want more information about the biodiversity value of their landholdings

#### 1.3 Limitations

The following limitations must be considered during the application of the bioregional plan:

- The bioregional plan does not replace the need for site assessments. Although the systematic biodiversity plan, which underpins the bioregional plan, was developed at a fine-scale, **on-site verification of the map of Critical Biodiversity Areas is always required**. The findings of site assessments must be interpreted based on the associated guidelines for the categories of Critical Biodiversity Areas;
- The systematic biodiversity plan (the Conservation Assessment and Plan for the Nelson Mandela Bay Municipality), which underpins the bioregional plan, represents the best available knowledge at the time of its development knowledge of some aspects of biodiversity, for example distribution of species of special concern, is not complete. In addition, our understanding of the

STEW/reeb Draft Bioregional Plan 2 Dec 10 for gazetting (2014-final)

Page 13

December 2010

biophysical features that are required to maintain essential ecological processes and to support the continued provision of important environmental goods and services continues to develop; and

• The landscape is constantly changing – significant changes in the landscape, for example large-scale loss of natural habitat, may impact on the location of areas that are required for conservation, and subsequently on the future configuration of the network of Critical Biodiversity Areas upon revision of the bioregional plan.


None of these limitations pose significant constraints to the application of the bioregional plan.

# 2 Part B: Biodiversity Profile

#### 2.1 Significance of the Biodiversity within the Bioregion

The global significance of the biodiversity of the Nelson Mandela Bay municipal area is emphasized by the fact that the area falls within two recognised biodiversity hotspots<sup>2</sup> (Myers *et al.*, 2000; Mittermeier, 2004). Biodiversity hotspots are the earth's richest and most threatened reservoirs of plant and animal life.

The rich biological diversity of the Nelson Mandela Bay is partly attributed to the fact that it is an area of convergence of five of South Africa's nine biomes<sup>3</sup> (see Figure 2-2): the Fynbos, Subtropical Thicket, Forest, Nama Karoo, and Grassland biomes (Low & Rebelo, 1998). Such a juxtaposition of biomes within a metropolitan municipality is unparalleled in the world (CI, 2009).



-

<sup>&</sup>lt;sup>2</sup> The Cape Floristic Region and the Maputaland-Pondoland-Albany Centre of Endemism

<sup>&</sup>lt;sup>3</sup> A biome is a regional-scale ecosystem characterized by distinct types of vegetation, animals, and microbes that have developed under specific soil and climatic conditions.

Page 14

Figure 2-1: Biodiversity hotspots of the world (Conservation International, 2004)




Figure 2-2: The biomes within Nelson Mandela Bay municipal area (Stewart *et al.*, 2008)

#### 2.2 Important Terrestrial Features

The NMBM Conservation Assessment identified a total of 58 vegetation types within the municipal area (see Table 2-1 and Figure 2-3).

STEW/reeb

Draft Bioregional Plan 2 Dec 10 for gazetting (2014-final)

Page 15




Figure 2-3: The historical distribution of natural vegetation types in the Nelson Mandela Bay municipal area (Stewart *et al.*, 2008)

Table 2-1: Key to Figure 2-3 – vegetation type numbers

No.	Vegetation Type	No.	Vegetation Type	No.	Vegetation Type
1	Albany Dune Thicket	21	Groendal Fynbos Thicket	41	Sundays Doringveld Thicket
2	Algoa Dune Thicket	22	Humewood Dune Fynbos	42	Sundays River
3	Baakens Forest Thicket	23	Intermediate Beach	43	Sundays River Floodplain
4	Baakens Grassy Fynbos	24	Koedoeskloof Karroid Thicket	44	Sundays Spekboom Thicket
5	Baviaans Spekboom Thicket	25	Kragga Kamma Indian Ocean Forest	45	Sundays Thicket
6	Bethelsdorp Bontveld	26	Lady Slipper Mountain Fynbos	46	Sundays Valley Thicket
7	Bushy Park Indian Ocean Forest	27	Lorraine Transitional Grassy Fynbos	47	Swartkops Escarpment Valley Thicket
8	Cape Recife Bypass Dunefield	28	Maitlands Dunefield	48	Swartkops Estuarine Floodplain
9	Chelsea Forest Thicket Mosaic	29	Malabar Grassy Fynbos	49	Swartkops Estuary
10	Coastal	30	Motherwell Karroid Thicket	50	Swartkops River
11	Coastal Hummock Dunes	31	Pan	51	Swartkops River Floodplain

STEW/reeb Draft Bioregional Plan 2 Dec 10 for gazetting (2014-final) December 2010

Page 16

December 2010

No.	Vegetation Type	No.	Vegetation Type	No.	Vegetation Type
12	Coega Estuary	32	Rocklands Renoster Bontveld	52	Swartkops Salt Marsh
13	Coega Estuary Floodplain	33	Rocklands Valley Thicket	53	Thornhill Forest and Thornveld
14	Colchester Strandveld	34	Rocky Beach	54	Van Stadens Afro. Ind. Ocean Forest
15	Colleen Glen Grassy Fynbos	35	Rowallan Park Grassy fynbos	55	Van Stadens Forest Thicket
16	Driftsands Bypass Dunefield	36	Sandy Beach	56	Van Stadens River
17	Driftsands Dune fynbos	37	Sardinia Bay Forest Thicket	57	Walmer Grassy Fynbos
18	Goudini Grassy Fynbos	38	Schoenmakerskop Rocky Shelf Fynbos	58	Wetland
19	Grass Ridge Bontveld	39	Skurweberg Grassy Fynbos		
20	Groendal Fynbos	40	St Francis Dune Fynbos Thicket Mosaic		

A high proportion of the vegetation types within the municipal area are threatened (twenty-three are *Critically Endangered* and a further ten are *Endangered*)<sup>4</sup> and now face collapse unless efforts are made to ensure their persistence (see Table 2-2 and Figure 2-4).

Table 2-2: An overview of the Ecosystem Threat Status of the vegetation types of the NMBM area

Ecosystem Threat Status	Number of Vegetation Types
Critically Endangered	23
Endangered	10
Vulnerable	15
Least Threatened	10
Total	58

The threatened status of these habitats is mirrored by the number of threatened species within the municipal area – the Nelson Mandela Bay Municipality has the highest number of red listed species of all the municipalities in the Eastern Cape (CSIR, 2004).

The most threatened vegetation types within the NMBM area are Lowland Fynbos, Subtropical Thicket mosaic and Forest vegetation types.

The Lowland Fynbos vegetation types in the NMBM area are predominantly grassy fynbos systems. Due to the fact that much of the urban expanse of Port Elizabeth was developed on these ecosystems, they are also some of the most threatened vegetation types within the municipal area. The most threatened Lowland Fynbos vegetation types within the municipal area are *Critically Endangered* Colleen Glen Grassy Fynbos, Lorraine Transitional Grassy Fynbos, Baakens Grassy Fynbos and Walmer Grassy Fynbos. Threatened plant species in these systems include South Africa's national flower, the king protea *Protea cynaroides* (*Critically Endangered*), *Cyclopia pubescens* (*Critically Endangered*) and *Brunsvigia litoralis* (*Endangered*). Pressures on the Lowland Fynbos within the NMBM area include invasion by alien vegetation, urban expansion, inappropriate fire regimes, habitat fragmentation and global climate change (Pierce-Cowling, 2009).

-

<sup>&</sup>lt;sup>4</sup> The Ecosystem Threat Status of the vegetation types in the Nelson Mandela Bay municipal area was assessed using biodiversity targets, which were based on the species-area curve method (Desmet & Cowling, 2004). The Ecosystem Threat Status of the vegetation types differs from the Draft National List of Threatened Ecosystems in Notice 1477 of 2009 in Government Gazette No. 32689 dated 6 November 2009. This difference is due to the fine-scale spatial mapping of the NMBM vegetation types, as opposed to broader level mapping that informed the national listing.

Page 17




Figure 2-4: The Ecosystem Threat Status of vegetation types within the Nelson Mandela Bay municipal area (Stewart et al., 2008)

The most threatened Subtropical Thicket vegetation types occurring within the municipal area are those that form mosaics with vegetation that have affinities to other biomes and include *Endangered* Sundays Doringveld Thicket, Bethelsdorp Bontveld and Motherwell Karroid Thicket. Coastal Fynbos Thicket mosaics within the municipal area are confined to the southern coastline and are a mixture of fynbos and large-leaved subtropical shrubs. They include *Critically Endangered* Humewood Dune Fynbos and *Endangered* Schoenmakerskop Rocky Shelf Fynbos, Driftsands Bypass Dunefield, Driftsands Dune Fynbos and St Francis Dune Fynbos Thicket Mosaic. Threatened plant species in the Subtropical Thicket mosaics include *Critically Endangered Aspalathus recurvispina*, *Orthopterum* 

STEW/reeb

Draft Bioregional Plan 2 Dec 10 for gazetting (2014-final)

December 2010.

Page 18

coegana and Aloe bowiea. The Albany adder Bitis albanica (Critically Endangered) also occurs in thicket and thicket mosaic habitat types. The mosaics are threatened by the development of informal and formal housing, industrial development, plant harvesting for medicinal and cultural use, overgrazing and mining.

Forest in Nelson Mandela Bay occurs in a solid form and in a mosaic with thicket. The solid forest within the NMBM is mostly confined to steep riverine gorges, which are sheltered from fire, and confined to a band along the wetter southern portion of the municipal area. Threatened forest types within the municipal area include *Critically Endangered* Chelsea Forest Thicket Mosaic, Bushy Park Indian Ocean Forest, Van Stadens Afromontane Indian Ocean Forest, Van Stadens Forest Thicket, Kragga Kamma Indian Ocean Forest and Thornhill Forest and Thornveld. The protected milkwood *Sideroxylon inerme*, dwarf Cape Beach *Rapanea gilliana*, and yellowwood *Afrocarpus falcatus* trees are prominent in many of the forest types. The southern population of blue duiker Philantomba monticola, which is confined to the Maputaland-Pondoland-Albany hotspot, occurs in the southern forested portions of Nelson Mandela Bay. The forest within the municipality is threatened by increased prevalence of fire, harvesting of forest products and the development of informal and formal housing

#### 2.3 Important Freshwater Features

Nelson Mandela Bay has ten main rivers, each with a unique ecosystem (Roux *et al.*, 2002), of which four are deemed to no longer be intact (SRK Consulting, 2009). Since all of the main rivers in Nelson Mandela Bay are unique, they all require some degree of safe-guarding.

Of particular importance is the Swartkops River system. The Swartkops River catchment contains two main rivers systems, the Elands and the Kwazunga, and two subsidiary tributaries, the Brak and Chatty. The Groendal Wilderness Area constitutes almost a third of the catchment of this river system, and the Groendal Dam supplies almost two-thirds of Uitenhage's domestic water. A number of threatened and rare species occur in Groendal, including Hewitt's ghost frog *Heleophryne hewitti* (*Critically Endangered*) and the Cape redfin minnow *Pseudobarbus afer*, which is restricted to the area.

Cultivation and invasive alien plants along this river system have impacted on its functioning by respectively increasing sedimentation and reducing the volume of water within the river. Alien species, such as the largemouth bass *Micropterus salmoides*, are a threat to the survival of indigenous fish species.

A number of small views and pans occur within the municipal area, particularly in grassy Fynbos, Subtropical Thicket, and coastal areas. These provide distinct habitats for species, including protected amphibian species such as the African bullfrog *Pyxicephalus adspersus* and the yellow-striped reed frog *Hyperolius semidiscus*. In addition to providing habitat, these wetlands are also important for the functions that they provide, such as flood attenuation, water filtration and erosion reduction. The main pressures on wetlands are through infilling or draining (e.g. for urban development and grazing), erosion, water abstraction and diversion (e.g. through dams and weirs).

#### 2.4 Important Coastal Features and Estuaries

The coastal zone of Nelson Mandela Bay provides an array of recreational and sustainable resource utilisation opportunities and underpins a substantial segment of the municipality's economy.

STEW/reeb Draft Bioregional Plan 2 Dec 10 for gazetting (2014-final) December 2010

Page 19

The Swartkops Estuary lies about 8 km north of the Port Elizabeth city centre and is an important nursery for marine fish, including several angling species. The estuary forms the basis of an informal bait collecting industry and is ranked as South Africa's top temperate estuary in terms of subsistence value (Turpie & Clark, 2007). The bait collecting industry is also, however, responsible for destruction of the mudflats, leading to significant impacts on the functioning of the system (Baird *et al.*, 1988). The functioning of the estuary is also threatened by pollution, enrichment, sand mining, overfishing, insufficient inflow of freshwater due to damming, and inappropriate development.

The foredune vegetation along the Nelson Mandela Bay coastline helps to buffer against the high energy influences of the tides, wind and waves. These systems are sensitive to disturbance and are susceptible to the formation of blow-outs. The coastal area is often targeted for development, primarily due to the desirability of ocean views. The protection of the coastal foredune system is important in order to safeguard the scenic attributes of the coastal zone and to provide protection against the natural coastal agents of change, particularly in light of the implications of global sea-level rise.

The Alexandria Dunefield has its beginnings at the north-eastern extent of the municipal area. The dunefield is considered to be one of the largest and most pristine active coastal dunefields in the world. Larger and more extensive dunefields are only found in desert areas. As such, it has been incorporated into the Addo Elephant National Park and was nominated by South African National Parks as a World Heritage Site (UNESCO, 2009).

The Maitlands Dunefield, between the resort villages of Beachview and Blue Horizon Bay on the south-western part of the Nelson Mandela Bay coastline, supports what is believed to be the highest density of black oystercatcher in the country (Bornman & Klages, 2003). The black oystercatcher is *Near Threatened* and is highly susceptible to disturbance during its breeding cycle. The Maitland area, in general, is one of the most visually spectacular areas in the municipality and has significant value in terms of recreation and tourism.

Although the prohibition of off-road vehicles on the dunefields has substantially reduced threats to these systems, they continue to be threatened principally by encroachment of invasive alien plants and are vulnerable to insensitive recreational use.

#### 2.5 Important Ecological Corridors

A number of ecological corridors were identified as part of the NMBM conservation assessment. These corridors were designed to promote connectivity between natural areas in order for ecological processes (such as migration and seed dispersal) to continue.

While all of the identified corridors are important, the Baakens River Valley, which forms an east-west corridor through the urban expanse of the municipal area, deserves a special mention. It is the municipality's most extensive corridor through fynbos habitats and is of critical importance for the continuation of ecological processes that sustain biodiversity. The area also provides numerous ecosystem services, playing an important role in flood attenuation, storm water management, environmental education and nature-based recreation.

#### 2.6 Patterns of Land-Use

A large proportion (62%) of Nelson Mandela Bay is in a natural state (SRK Consulting, 2009), making it the metropolitan municipality with the highest proportion of natural habitat in the country.

STEW/reeb Draft Bioregional Plan 2 Dec 10 for gazetting (2014-final)

Page 20




Figure 2-5: Patterns of land-use in the Nelson Mandela Bay municipal area (SRK Consulting, 2009)

This creates opportunities that most metropolitan municipalities no longer have for capitalising on ecological infrastructure and for natural ecosystems to support human well-being (e.g. through ecosystem-based adaptation to climate change).

Urban development is primarily concentrated on the coastal and inland lowlands in the east and northwest of the municipal area. Cultivation is concentrated on the ancient dunes in the south of Nelson Mandela Bay and the inland Fynbos and Subtropical Thicket lowlands (see Figure 2-5).

STEW/reet

From 2003 development within the coastal zone has increased substantially. The extent of habitat loss within the coastal zone is of concern not only due to impacts on ecological processes and biodiversity but also due to the impacts on the natural character of this zone, which is a key component of the municipality's tourism economy (SRK Consulting, 2009).

# 2.7 Ecosystem Protection Level within the Nelson Mandela Bay Municipal Area

The existing network of conservation areas within the NMBM area totals approximately 10,500 ha, or approximately 5 % of the municipal area. Of this, only 4,700 ha (2 % of the municipal area) is recognised as protected areas in terms of the NEM: Protected Areas Act (Act No. 57 of 2003). The remaining conservation areas (approximately 5,800 ha in extent) have weak legal protection and are not necessarily managed for biodiversity conservation purposes.

The current protected area system in the municipal area is therefore highly deficient, particularly in terms of its limited size and connectivity, and urgently requires expansion in order to achieve biodiversity targets (SRK Consulting, 2009). Due to the extent of remaining natural habitat, the NMBM is unique amongst the metropolitan municipalities in terms of opportunities for expanding the protected area network to the benefit of biodiversity as well as its people.

Table 2-3: Distribution of conservation areas within the municipal area

Protected Areas	Area (ha)
Greater Addo Elephant National Park	752
Groendal Wilderness Area	14
Island Forest Nature Reserve	496
Cape Recife Nature Reserve	323
Lady Slipper Nature Reserve	365
Maitland Nature Reserve	132
Sardinia Bay Nature Reserve	249
Settlers Park Nature Reserve	75
Swartkops Valley Nature Reserve	940
Sylvic Nature Reserve	91
The Springs Nature Reserve	929
Van Stadens Nature Reserve	339
Total	4,705

Conservation Areas	Area (ha)
Grassridge Private Nature Reserve	1,320
NMMU Private Nature Reserve	840
Witteklip Private Nature Reserve	29
Van Stadensberg Natural Heritage Site	579
Seaview Game Park	31
Kragga Kamma Game Park	203
Paardekop Game Farm	679
Tregathlyn Game Farm	2,099
Total	5,780

#### 2.8 Description of Critical Biodiversity Areas

Critical Biodiversity Areas within the municipal area are the portfolio of sites that are required to meet the municipality's biodiversity targets<sup>5</sup>. Such areas should be maintained in their natural state in perpetuity. A map of CBAs was produced as part of the NMBM's conservation assessment and sites

STEW/reeb Draft Bioregional Plan 2 Dec 10 for gazetting (2014-final) December 2010

<sup>&</sup>lt;sup>5</sup> The biodiversity targets for each of the vegetation types in the municipal area were assigned according to the targets for corresponding broad habitat units or vegetation types as identified in the C.A.P.E. and STEP Programmes.

Page 22

were assigned to CBA categories based on their biodiversity characteristics and Ecosystem Threat Status (see Table 2-4).

Table 2-4: Description of the Critical Biodiversity Area categories

Category	Code	Description	Land Management Objective
Protected Area 1	PA 1	Protected areas managed by SAN Parks, provincial or local authorities, parastatals (e.g. NMMU), or the private sector. Includes National Parks, Provincial, Local and Private Nature Reserves.	To be maintained as Protected Areas.
Protected Area 2	PA 2	National Parks, Provincial, Local, Private Nature Reserves pending declaration.	To be declared and maintained as Protected Areas.
Critical Blodiversity Areas	ÇBAs	All Critically Endangered habitats, ecological process areas, ecological corridors, habitats for Species of Special Concern, and some Endangered, Vulnerable or Least Threatened habitats.	Such areas must be managed for biodiversity conservation purposes and incorporated into the protected area system.
Ecological Support Area 1	EŜA 1	Agricultural land that plays an important role in ecosystem functioning and I or provides connectivity between natural areas.	Such areas must be maintained for extensive agricultural purposed and managed to promote ecological connectivity.
Ecological Support Area 2	ESA 2	Areas severely disturbed or transformed by human activities (e.g. mining), requiring restoration or rehabilitation.	Such areas must be restored or rehabilitated to support ecological connectivity. Such areas must not be developed.
Other Natural Areas	ONA	Natural areas that are not required to meet biodiversity targets.	As per the Municipal SDF or local SDFs.
Areas where no natural habitat remains	DEV	Areas severely disturbed or transformed by human activities with no natural habitat remaining, including airfields, cultivated lands, forestry plantations, industry, mines and quarries, severe overgrazing, and urban and rural development.	As per the Municipal SDF or local SDFs.

The CBAs were clustered into twenty-nine implementation sites (see Figure 2-6) and were prioritised according to their overall Ecosystem Threat Status and current land-use. These implementation areas should form the focus for the investment of the limited resources of the NMBM for the implementation of the bioregional plan.

Page 23




Figure 2-6: Implementation sites in the Nelson Mandela Bay municipal area (SRK Consulting, 2009)

Page 24

# 3 Part C: Critical Biodiversity Areas map




Figure 3-1: Map of Critical Biodiversity Areas within the municipal area

# 4 Part D: Guidelines

Table 4-1: Biodiversity-compatible land-use guidelines matrix - Category descriptions and management objectives<sup>6</sup>

Category on the CBA Map	Description	Land Management Objective	Land Management Recommendations	General Land-Use	Associated NMBM Rural Development Policy and SDF Categories
Protected Areas (PA 1 & 2)	Formal Protected Areas (PA1) and Protected Areas pending declaration (PA2), in terms of the National Protected Areas Act and ECA.	Maintain or restore NATURAL structure and ecosystem functioning.	Maintain formal conservation protection.	Conservation (non consumptive).	Critical Biodiversity Zone 1 (CBZ1)
Critical Biodiversity Areas (CBAs)	ANY land or aquatic site required to meet biodiversity pattern (features), process (ecological), or species targets.	Maintain or restore NATURAL structure and ecosystem functioning.	Maintain or obtain formal conservation protection.	Conservation (non consumptive).	Critical Biodiversity Zone 1 or 2 (CBZ1 or 2)
Ecological Support Area 1 (ESA1)	Agricultural land essential for connectivity in the landscape, particularly between CBAs.	Maintain existing extensive land-uses (e.g. sustainable livestock grazing) where these facilitate ecological functioning and connectivity between adjacent CBAs; or restore NATURAL structure and ecosystem functioning; or rehabilitate to near natural state to facilitate connectivity between adjacent CBAs.	Maintain existing extensive landuses or manage for conservation.	Extensive agriculture that promotes conservation (sustainable consumptive or non consumptive).	Critical Biodiversity Zone 2 (CBZ2)
Ecological Support Area 2 (ESA2)	Severely disturbed or degraded sites (e.g. mines, overgrazed areas) essential for ecosystem connectivity in the landscape, particularly between CBAs.	Restore NATURAL structure and ecosystem functioning, or rehabilitate to facilitate connectivity between adjacent CBAs.	Obtain formal protection and manage for conservation once existing land-use has ceased.	Conservation (sustainable consumptive or non-consumptive) once existing landuses causing degradation have ceased.	Critical Biodiversity Zone 2 (CBZ2)
Developed Areas: Non-restorable land with no natural habitat remaining (Dev)	Irreversibly developed / degraded areas (areas that cannot be rehabilitated):  Intensive agriculture (excluding ESA1s);  Urban development;  Industry; and  Waste sites etc.	As per Municipal SDF, local SDF and NMBM Urban Densification and Rural Development Policy.	As per Municipal SDF, local SDF and NMBM Urban Densification and Rural Development Policy.	Preferred sites for development, but always subject to conditions in terms of NMBM SDF, Urban Densification and Rural Development Policy and 'sustainable human settlement' principles.	Various

<sup>&</sup>lt;sup>6</sup> With acknowledgement to Deborah Vromans (Garden Route Initiative [GRI], SAN Parks) who compiled initial draft guidelines for the Garden Route Initiative, which were used as a template during the development of these guidelines.

Draft Bioregional Plan 2 Dec 10 for gazetting (2014-final)

Page 26

SRK Consulting Draft Bioregional Plan

Associated NMBM Rural Development Policy and SDF Categories	Various
General Land-Use	Preferred sites for development, but always subject to conditions in terms of NMBM SDF, Urban Densification and Rural Development Policy and 'sustainable human settlement' principles.
Land Management Recommendations	As per Municipal SDF, local SDF and NMBM Urban Densification and Rural Development Policy.
Land Management Objective	As per Municipal SDF, local SDF and NMBM Urban Densification and Rural Development Policy.
Description	Natural & intact but not required to meet targets.
Category on the CBA Map	Other Natural Areas (ONA) (TERRESTRIAL and AQUATIC) Including degraded areas (e.g. high density alien plants) not part of CBAs and ESA 1 & 2.

Draft Bioregional Plan 2 Dec 10 for gazetting (2014-final)

Page 27

# Table 4-2: Recommended land & resource (water) use management guidelines in Critical Biodiversity Areas and other categories on the CBA map

#### **CRITICAL BIODIVERSTIY AREAS (TERRESTRIAL ECOSYSTEMS)**

NCLUDES: ALL Critically Endangered vegetation types in terms of the NMBM Conservation Plan and Best Design sites of Endangered, Vulnerable & Least Threatened vegetation types, ecological processes and species of special concern that satisfy biodiversity tarriets

#### Keep in a NATURAL STATE

#### Biodiversity Friendly Land-Use Types (Land-Use Type 1-4)

#### OBJANCANIER N

#### **General Recommendations**

- No further loss of natural habitat should occur i.e. all land in this category should be maintained as natural vegetation cover;
- These areas of land can act as possible biodiversity offset receiving areas;
- Mimic or maintain natural ecological processes; for example fire in Fynbos and Dune Thicket vegetation types;
- Determine the vegetation type and then consult the Fynbos Forum Ecosystem Guidelines (de Villiers et al., 2005) for detailed management guidelines;
- Priority for DEDEA, NMBM, Department of Agriculture, and Forestry Section of DAFF to carry out regular environmental
  monitoring and reporting, for biodiversity and / or change of land-use, to prevent unauthorized development or degradation by
  neglect or ignorance (monitoring guidelines are required);
- Prioritise CBAs for land care projects, Working for Water (WfW) and NGOs to direct their conservation projects, programmes and activities; and
- An Ecological Management Plan should be compiled by landowner for CBAs. EMP to include alien plant control, fire management etc.

#### **Protection**

CBAs not formally protected (public or private) should be rezoned to Open Space 3, declared in terms of NEM: Protected Areas
Act and privately owned erven to be incorporated into the protected area network through Conservation Agreements and
incentives. Title deed restrictions must be attached to these. Rates rebates can be provided in return.

#### Fire Management

• An appropriate fire management plan for fire dependant ecosystems must be implemented, namely Fynbos and Dune Thicket.

#### Rehabilitation / Restoration

 Degraded or disturbed CBAs will require restoration or rehabilitation through programmes such as Working for Water, Working for Wetlands.

#### In Urban Areas

- Maintain and legally protect the Urban Open Space System that maintains CBAs; and
- Compatible land-uses adjacent to and within CBAs & management guidelines could include
  - Community projects such as passive recreation for eco-tourism, employment of security officers (visible 'policing') etc.
  - Where necessary, provide fencing or locate institutional / educational 'barriers' where high density, high canopy vegetation types create a security risk.

#### Sand Movement (Ecological) Corridors

- No development should be allowed in sand movement corridors (littoral active zone), frontal dunes or dynamic dune systems;
- Mobile dunes must not be stabilized, unless where existing development requires it;
- Maintain an indigenous vegetation buffer zone between the inland boundary of the youngest fixed dune trough and the seaward boundary of existing development; and
- Strict control should be maintained over the use of off-road vehicles on beaches.

#### **Coastal Ecological Corridors**

Due to expected sea level rise from climate change, no development should occur within the coastal CBAs and within 300 m of
the high water mark outside of the urban edge, or within CBAs and below 10 meters above sea level within the urban edge.

STEW/reeb

Draft Bioregional Plan 2 Dec 10 for gazetting (2014-final)

Page 28

#### **Development Guidelines**

#### Where infrastructure is proposed, the following guidelines should be implemented -

- Rezoning of properties to afford additional land-use rights that will result in increased biodiversity loss should not be granted;
- Permitted number of units per erf or per ha to be based on existing land-use rights and NMBM Rural Development and Urban Edge Policy;
- Developments should be limited to existing developed / degraded footprints, if present;
- Units carefully dispersed or clumped to achieve least impact, particularly with regard to habitat loss and fragmentation;
- The following minimum buffers are recommended between CBAs and proposed development nodes -
  - Forest: outside urban edge: minimum of 100 m, urban areas minimum of 30 m; and
  - Other biomes: minimum of 50 m;
- The installation of infrastructure in CBAs is not desirable and should only be considered if all alternative alignment and design
  options have been assessed and found to be non-viable. Under such conditions, at least a Basic Assessment (BA) should be
  undertaken, and if approved, a comprehensive EMP must be developed and best-practice restoration efforts strictly
  implemented;
- Protected Area infrastructure to be limited to existing developed or degraded sites, unless determined otherwise via a Conservation Development Framework planning process for the reserve;
- Ecological Specialist to conduct the ecological assessment;
- Where the infrastructure is within close proximity to aquatic ecosystems refer to 'CBA Aquatic Ecosystems' below to ensure appropriate buffers;
- Temporary structures are preferred e.g. raised wooden structures, tents, raised boardwalks or alternatively porous materials / design concepts preferred; and
- A Construction & Operational Environmental Management Plan to be compiled and implemented.

Where development proposals other than the preferred biodiversity-compatible land-uses (see definitions below in Table 4-3, Table 4-4, Table 4-5 and Table 4-6) are submitted in terms of the NEMA: EIA regulations or Land Use Planning Ordinance (LUPO):

- · A Screening Exercise should be undertaken by a Biodiversity Specialist/Ecologist to verify the CBA map category on site;
- If the site is verified as a CBA, developments other than Conservation Use should be identified as inappropriate;
- · If the application is pursued they should be informed by a specialist biodiversity assessment; and
- Developments outside of the Urban Edge are also subject to the NMBM Rural Development Policy.

Page 29

#### **CRITICAL BIODIVERSITY AREAS (AQUATIC ECOSYSTEMS)**

\*Critically Endangered and Critical Ecological Process sites – in terms of NMBM pattern & ecological process targets INCLUDES wetland types, river reaches, estuaries and catchments

#### Keep in a NATURAL or NEAR NATURAL state

Conservation Resource Use – HIGH resource PROTECTION

Adjacent Land-Use – CONSERVATION or buffer to adjacent development

#### **BIODIVERSITY OFFSETS**

No realistic biodiversity offsets.

#### **ECOLOGICAL RESERVE**

#### **Water Quantity**

- · Maintain mean annual run-off as close to natural as possible;
- Where abstraction is absolutely necessary, off channel dam construction ONLY with the highest Ecological Reserve possible;
- Improve regulation of abstractions from these rivers & encourage efficient water use throughout the area (i.e. demand control, recycling, use of grey water, rainwater tanks); and
- Where water abstraction is licensed, all effluent must be treated and returned to natural water courses in order to obtain maximum utilization of scarce water resources.

#### **Environmental Flow Requirements**

- Flow regimes as close to natural as possible Retaining natural flow regimes (both in terms of magnitude and variability);
   Management actions to maintain natural flow regime should include:
  - Abstraction should occur during high flow seasons, not during low flow seasons;
  - Water release from dams should take note of the ecological requirements of CBA rivers. This includes at least one annual flood release (at most appropriate time of year for each system), even if the dam is not full; and
  - Optimal use should be made of existing abstractions through demand-management measures. Water must be correctly
    priced to prevent abuse and wastage.

#### **Water Quality**

- Water quality as close to NATURAL as possible;
- The Target Water Quality Range (TWQR) (range of concentrations or levels) should ensure that no measurable adverse effects are expected on the health of aquatic ecosystems, and should therefore ensure their protection;
- All effluent is treated and returned to natural water resources in order to obtain maximum utilization of scarce water resources (as per DWA Policy); and
- A precautionary approach is required to protect the health of aquatic ecosystems, which means that active measures are taken
  to avert or minimise potential risks of undesirable impacts on the environment.

#### **Monitoring & Management**

- A management plan should developed for each biodiversity feature or site, for example red data species, specific river reach habitat and
- NMBM to investigate establishment of aquatic reserves (e.g. Swartkops Estuary).

#### Buffer Recommendations (CBA Buffers are 'NO-GO' areas)

Where an aquatic CBA is situated within a Ecological Corridor CBA, the width of the CORRIDOR will apply in concert with the buffers recommended below. UNLESS other land-uses (e.g. Agriculture) exist already OR a conservation land-use is proposed, the buffers recommended below will apply & should be actively enforced.

It is important to NOTE that the following buffer recommendations DO NOT REPLACE a SITE LEVEL investigation.

STEW/reeb

Draft Bioregional Plan 2 Dec 10 for gazetting (2014-final)

Page 30

#### **River Buffers**

To protect river integrity the following general buffers will apply:

- Mountain streams and upper foothills of all 1:500,000 (major) rivers = 50 m (these rivers generally have more confined riparian
  zones than lower foothills and lowland rivers; and are generally less threatened by agricultural practices);
- Lower foothills and lowland rivers of all 1:500,000 rivers = 100 m (these rivers generally have less confined riparian zones than
  mountain streams and upper foothills; and are more threatened by agricultural practices. These larger buffers are particularly
  important to lower the amount of pesticides etc. reaching the river); and
- All remaining 1:50,000 **streams** = 32 m (these rivers are generally the smaller upland streams corresponding to mountain streams and upper foothills. They are generally smaller rivers than those designated in the 1:500,000 rivers layer, and are assigned the riparian buffer required under South African legislation).

#### Alternatively

- If the 1:100 year flood line (or if flooding has occurred > 1:100 year flood line) is greater than the buffers prescribed above, then
  these flood lines will apply (i.e. it is important to re-evaluate flood lines to cater for the impact of climate change and reduce
  flood damage on new developments); and
- Enforcing the above riparian buffer zones along aquatic CBAs is crucial. The buffers will apply to crops, since rivers and their
  associated biota are highly susceptible to crop pesticides. It also applies to excluding livestock, which cause considerable bank
  erosion, with subsequent degradation of water quality. The access areas for livestock should be demarcated and all alien
  invasive plants within this zone should also be eradicated.

#### **Estuary Buffer**

- It is recommended that no new development around an estuary be permitted below 500 m from the high water mark or the 1:100 year flood line, whichever is the greatest;
- Where known flooding has occurred (e.g. Swartkops River Estuary) the highest flood line will apply (i.e. re-evaluate flood lines
  to cater for the impact of climate change and reduce flood damage on new developments); and
- Estuarine salt marshes should be protected and no activities allowed there.

#### **Wetland Buffers**

- Wetlands classified as CBA and that are largely intact (natural to near natural), namely rank 1 & 2 wetlands, require a minimum buffer of 75 - 200 m;
- Ensure that delineation of the wetland boundary is undertaken by a specialist ecologist according to 'A practical field procedure for identification and delineation of wetlands and riparian areas' (DWAF, 2003);
- No roads should be permitted to traverse these wetlands & their buffers; and
- Where a road or other water channelling structure runs close to a wetland and its buffer, and channels water into it, such water should be dispersed via multiple entry points with energy-dispersing structures. These drains must be small, dispersed low-volume, low-velocity structures. They must also be set back from the wetland and its buffer zone and be designed to spill into undisturbed natural vegetation at ground level. These provisions also apply to all urban storm-water outlets that spill into a wetland or up-slope of a wetland.

The following buffers are recommended for rank 1 and 2 wetlands, which vary according to size and wetland integrity (rank):

Size of Wetland	Rank 1	Rank 2	
>20 ha	200 m	150 m	
5 – 20 ha	150 m	100 m	
< 5 ha	100 m	75 m	

#### **General Recommendations regarding Buffers -**

Page 31

- Do not permit infilling, excavation, drainage, hardened surfaces (including buildings and asphalt), intensive agriculture or any
  new developments within a river, wetland or estuary, their buffers, the 1:100 year flood line or highest flood line of the
  delineated edge, whichever is the greatest.
- The "buffers" may need to be wider than the above recommendations, for example due to the presence of an important amphibian species. This should be determined on a case-by-case basis by a specialist ecologist in consultation with the Department of Water Affairs (DWA), the NMBM and DEDEA to reflect site-specific factors. The approach for determining buffer width should consider the current condition of the aquatic ecosystem, as well as the functioning of the system in the broader landscape, plus an assessment of the impacts to the ecosystem of the existing and proposed adjacent land-use and climate change impacts i.e. increased flooding at higher levels than previously recorded.

#### REHABILITATION

- Areas that are degraded or disturbed should be rehabilitated, through programmes such as Working for Wetlands and a systematic alien vegetation eradication programme implemented to improve biodiversity and water supply, especially upstream areas of estuaries and wetlands;
- Prohibit the stocking of farm dams (even off-stream dams) with alien fish;
- Remove alien fish where possible to allow for natural recovery of indigenous fish in consultation with DWA, DEDEA, EC Parks & NMBM;
- Stock dams with indigenous fish in consultation with DWA, DEDEA, EC Parks & NMBM; and
- Rehabilitate riparian zones to act as a buffer between the river and surrounding agricultural areas in consultation with the Department of Agriculture, Forestry and Fisheries (DAFF), Department of Water Affairs (DWA), DEDEA & NMBM.

#### MANAGING CUMULATIVE IMPACTS THROUGH TRANSFORMATION THRESHOLDS WITHIN CBA CATCHMENTS

- Development in CBA catchment areas should be biodiversity-friendly land-uses that discourage the following activities in upstream catchment areas to prevent unacceptable transformation levels & impacts on inland wetlands, estuaries and lowland rivers:
  - Large-scale abstraction, river diversion, impoundments, urban development, extensive woody alien invasion, river diversion / water transfer, Waste Water Treatment Works (WWTW) discharges, poorly serviced informal settlements, cultivation, impacts to floodplain hydrology, changes in water table (resulting from catchment-scale water transfers), catchment hardening.
- Catchment Developments (e.g. forestry) remain subject to standard DWA requirements regarding limits to reduction in mean annual runoff.

#### OTHER GENERAL MANAGEMENT GUIDELINES

- Aquaculture projects associated with aquatic CBAs e.g. off-stream dams, should not be permitted;
- Hydrological connections between systems (surface or groundwater) should not be disrupted (includes the need to manage post-development high flow and low flow runoff volumes);
- Manage jetties and structures on rivers and estuaries;
- Manage bio-physical and recreational carrying capacity of rivers and estuaries;
- Manage estuary mouth dynamics as close to natural as possible;
- Most of the estuaries & downstream wetlands are highly threatened by loss of freshwater inflows from upstream and by development encroachment. Management of these issues is considered critical;
- An Environmental Management Plan to be compiled for adjacent land-uses and should address the following issues: buffers, water quality, water flow, abstraction thresholds, alien fauna control, storm water etc.; and
- The Fynbos Forum Ecosystem Guidelines for Environmental Assessment in the Western Cape (de Villiers *et al.*, 2005) includes a section on river, estuary and wetland ecosystems that complements these guidelines.

Page 32

#### **ECOLOGICAL SUPPORT AREAS 1 (ESA1 - AGRICULTURAL LAND)**

Supporting zone required for maintaining connectivity between CBAs, continued ecosystem functioning within the CBA corridors, and preventing degradation of Critical Biodiversity Areas. These require high protection.

INCLUDES extensive agricultural land immediately adjacent to CBAs

Maintain agricultural land in an EXTENSIVE state or rehabilitate or restore to a natural or semi-natural state.

#### **General Recommendations**

- No further loss of natural habitat should occur and all land in this category currently in an extensive agricultural state should be
  maintained as such, or rehabilitated or restored to a natural or semi-natural state;
- The maintenance of connectivity between CBAs, continued ecosystem functioning within the CBA corridors, and the prevention
  of degradation of adjacent Critical Biodiversity Areas must be achieved;
- All natural ecological processes on adjacent CBAs to continue e.g. fire in Fynbos, Dune Thicket vegetation types;
- No agri-industry, urban residential, mining, or more intensive land-uses (e.g. cultivation, pastures etc.) other than the current state to be permitted;
- Development limitations as per the NMBM Rural Development Policy must be applied (e.g. max. subdivision size, max. residential density, maximum cluster footprint size etc.);
- Priority for DEDEA, NMBM & Department of Agriculture to carry out regular environmental monitoring and reporting, for biodiversity and / or change of land-use, to prevent unauthorized development or degradation by neglect or ignorance (monitoring guidelines are required);
- Prioritise these areas for land care projects, Working for Water, Working for Wetlands and NGOs to direct their rehabilitation / restoration projects, programmes and activities; and
- An Environmental Management Plan should be compiled, where possible, to include, e.g. alien plant control, fire management etc.

#### **Development Guidelines**

Where infrastructure is developed, the following guidelines should apply -

- Permitted number of units per erf or per ha to be based on existing land-use rights and NMBM Rural Development Policy and Urban Edge;
- Rezoning of properties to afford additional land-use rights that will result in increased biodiversity loss should not be granted,
  unless significant nett conservation gains can be achieved, ecosystem functioning and connectivity of Ecosystem Support
  Areas (ESAs) will not compromised, and biodiversity impacts with regard to species and habitats <u>outside of CBAs</u> are also of
  an acceptable significance;
- Developments should be limited to existing developed / degraded footprint, if present;
- Units carefully dispersed or clumped to achieve least impact;
- Protected Area infrastructure to be limited to existing developed or degraded sites, unless determined otherwise via a Conservation Development Process for the reserve;
- If the cadastre is located partially or entirely outside of a CBA / ESA, development should be located outside the CBA / ESA;
- Ecological Specialist to conduct the ecological assessment;
- Where the infrastructure is within close proximity to aquatic ecosystems refer to 'CBA Aquatic Ecosystems' below to ensure appropriate buffers;
- Temporary structures are preferred e.g. raised wooden structures, tents, raised boardwalks or alternatively porous materials / design concepts; and
- A Construction & Operational Environmental Management Programme to be compiled and implemented.

Page 33

#### **ECOLOGICAL RESERVE**

#### **Water Quantity**

Maintain mean annual run-off to ensure Eco-status depending on the feature or process being protected.

#### **Environmental Flow Requirements**

• Preferable to abstract during high flow seasons.

#### **Water Quality**

· Water quality to reflect the recommended Eco-status depending on the feature or process being protected.

#### Monitoring & Management

- Site level studies will need to be conducted to determine what Eco-Status is required to appropriately protect the relevant feature or process.
- As per CBA

#### **Buffer Recommendation**

Both the NEMA 32 m buffer and CARA 10 m buffer are legally enforceable.

#### **Wetland Buffers**

Inland wetland (excluding estuaries) buffers were determined according to size and wetland integrity, and are recommended as follows:

Wetland Size	Rank 3	Rank 4	Rank 5
>20ha	75m	<b>75</b> m	50m
5 – 20ha	75m	50m	50m
< 5ha	50m	50m	50m

#### Activities discouraged within the 1: 20 year flood line of inland wetlands:

 Creation of berms, roads, culverts, canalisation, channelisation, alien vegetation, impoundment, abstraction, well points, stormwater or other point source inflows, irrigation return flows, grazing / trampling, agriculture, golf courses, suburban gardens, artificial deepening, drainage, aquaculture, recreational activities, salt mining.

#### Activities discouraged within the 1: 100 year flood line of inland wetlands:

Intensive agriculture, intensive livestock grazing, sports-fields that require draining / infilling, market gardening (where nutrient
enrichment is an issue), creation of flood control berms, roads that interrupt flood flows, poorly serviced informal settlements,
alien vegetation, suburban gardens, golf course greens, catchment hardening, roads, culverts, raised or lowered water table.

#### **General Recommendations regarding buffers:**

- Do not permit infilling, excavation, drainage, hardened surfaces (including buildings and asphalt), intensive agriculture or any
  new developments within a river or wetland, their buffers, the 1: 100 year flood line or highest flood line of the delineated edge,
  whichever is the greatest; and
- The "buffers" may need to be wider than the above recommendations. This should be determined on a case-by-case basis by a specialist ecologist in consultation with the Department of Water Affairs (DWA), NMBM and DEDEA to reflect site-specific factors. The approach for determining buffer width should consider the current condition of the aquatic ecosystem and any existing or proposed buffer, as well as the functioning of the system in the broader landscape, plus an assessment of the impacts to the ecosystem of the existing and proposed adjacent land-use and climate change impacts i.e. increased flooding at higher flood lines.

STEW/reeb

Page 34

#### ECOLOGICAL SUPPORT AREAS 2 (ESA 2- IMPORTANT DEVELOPED / DEGRADED SITES)

Supporting zone required for maintaining connectivity between CBAs, continued ecosystem functioning within the CBA corridors, and preventing degradation of Critical Biodiversity Areas. These sites require high protection.

INCLUDES developed / degraded land immediately adjacent to CBAs that was or is presently utilised for intensive purposes (e.g. mining)

#### Manage for SUSTAINABLE DEVELOPMENT

Resource Use - HIGH resource PROTECTION

Land-Uses that maintain SUSTAINABLE development

#### **General Recommendations**

- No further loss of natural habitat should occur and all land in this category currently in a degraded / developed state must be
  rehabilitated or restored to a natural or semi-natural state once the current land-use has ceased (e.g. mining operation);
- Legal requirements for restoration / rehabilitation must be enforced by NMBM and relevant authority (e.g. DMR, DEDEA, DEA);
- The maintenance of connectivity between CBAs, continued ecosystem functioning within the CBA corridors, and the prevention
  of degradation of adjacent Critical Biodiversity Areas must be achieved;
- All natural ecological processes on adjacent CBAs to continue e.g. fire in Fynbos and Dune Thicket vegetation types.
- Priority for DEDEA, NMBM, Department of Mineral Resources (DMR) & Department of Agriculture to carry out regular
  environmental monitoring and reporting, for biodiversity and / or change of land-use, to prevent unauthorized development or
  degradation by neglect or ignorance;
- Prioritise these areas for land care projects, Working for Water, Working for Wetlands and NGOs to direct their rehabilitation / restoration projects, programmes and activities;
- An Environmental Management Plan should be developed and implemented, and should include the following aspects, amongst others: alien plant control, fire management etc.

#### **Development Guidelines**

Where infrastructure is developed, the following guidelines should apply:

- Permitted number of units per erf or per ha to be based on existing land-use rights and NMBM Rural Development Policy and Urban Edge Strategy;
- Rezoning of properties to afford additional land-use rights should not be granted, unless significant nett conservation gains can
  be achieved, ecosystem functioning and connectivity of ESA's will not be compromised, and biodiversity impacts with regard to
  species and habitats <u>outside of CBAs</u> are also of an acceptable significance;
- Developments should be limited to existing developed / degraded footprint, if present;
- Units carefully clustered or dispersed to achieve least impact;
- Other degraded areas to be restored / rehabilitated;
- Protected Area infrastructure to be limited to the most developed or degraded sites, unless determined otherwise via a Conservation Development Process for the reserve;
- If the cadastre is located partially or entirely outside of a CBA / ESA, development should be located outside the CBA / ESA.
- Ecological Specialist to conduct the ecological assessment;
- Where the infrastructure is within close proximity to aquatic ecosystems refer to 'CBA Aquatic Ecosystems' below to ensure appropriate buffers;
- Temporary structures are preferred e.g. raised wooden structures, tents, raised boardwalks or alternatively porous materials / design concepts; and
- A Construction & Operational Environmental Management Plan to be compiled and implemented.

#### **ECOLOGICAL RESERVE**

Water Quantity: Minimum Standards, Environmental Flow Requirements, Rehabilitation etc.

Recommendations: same as for ESA 1s

STEW/reeb

Draft Bioregional Plan 2 Dec 10 for gazetting (2014-final)

Table 4-3: Definitions of biodiversity-compatible land-use types

SRK Consulting Draft Bioregional Plan

Conservation  This is a land-use where conservation is the major objective:  Areas which are currently legally protected; and  Areas which are designated CBA but are not yet legally protected and should be, either through declaration or through a biodiversity stewardship agreement, where possible.  Appropriate uses:  Non-impact directly related activities such as research, outdoor recreation and environmental education;  Non-consumptive activities, for example, low-intensity recreation and eco-touring trails, bird watching and overnight	Open Space Zone 3	a. Conservation Management, which is the most biodiversity-compatible land-use specifically aimed at maintaining and enhancing biodiversity in Protected Areas. As a form of land-use it extends to include production where, for instance, game animals are hunted or captured. CM occurs on private land when legally binding, written formalities are in place to ensure such management extends beyond the scope of current ownership.  b. Low impact eco-tourism development on extensive areas of natural intact vegetation. Good management practices are implemented with small, low density footprints, determined via a Conservation Development Framework or similar process. Impacts are low due to size, technology (operational activities) and design concepts e.g. enviro-loos, temporary structures, green architecture, use of natural materials.
Sustainable consumptive activities, such as wildflower harvesting (for the medical, culinary or flower industry), subject to an Environmental Management Plan (EMP) demonstrating the sustainability of harvesting.  Land-use densities and footprints as per NMBM Rural Management and Urban Edge Policy, and NMBM integrated Zoning Scheme. Where existing infrastructure or a development footprint already exists, such areas should be used for development, but within sustainable limits and as the Policy.		Passive recreation, such as hiking, horse riding, bird watching, game watching etc. reduces the impact of this land-use. This land-use should be coupled with rezoning of the undeveloped areas to Open Space 3 and declaration as a Nature Reserve or Protected Environment in terms of the NEM Protected Areas Act.  c. Sustainable consumptive activities, which include the sustainable harvesting of natural resources, such as wildflower harvesting (for the medical, culinary or flower industry), fishing in riverine reserves & bee farming. These activities will require stringent management of harvesting volumes. It will require a specialist to conduct a study on carrying capacity and develop a management plan to ensure appropriate stocking rates, harvesting techniques & volumes are implemented & managed. The land-use must be based on strict management guidelines and sustainable resource use principles, as advised by a specialist or the conservation body (NMBM Environmental Services Sub-Directorate, ECPB or DEDEA).

Draft Bioregional Plan 2 Dec 10 for gazetting (2014-final)

SRK Consulting Draft Bioregional Plan

This land-use utilises extensive areas of natural veld. These areas need to be well managed to prevent negative impact e.g. determining veld carrying capacity, stocking rates & stocking with indigenous game. Any intensification or overdevelopment (e.g. for tourism) of these areas will increase the negative	impacts.		
Agriculture Zone AND Resort Zone			
Extensive Game Farming  Includes: Game production and related tourism activities on extensive land portions of natural land cover; sustainable commercial hunting along with other consumptive and non-consumptive use of wild natural resources.  Game reserve, including lodge and other accommodation infrastructure, with strict limits on the level of development considered acceptable.	Assumes: Minimum size criteria for economic and ecological sustainability are applied in terms of carrying capacity; strictly limited development for revenue generating purposes such as intensified tourism or sectional ownership.  Stringent management conditions applied, such as —	<ul> <li>Faunal specialist to undertake carrying capacity study;</li> <li>Indigenous or ecologically benign extra-limital species only;</li> <li>Environmental Management Plan;</li> <li>Fire management measures, if necessary; and</li> </ul>	<ul> <li>Avoidance of areas containing Critically Endangered or Endangered ecosystems and RDB listed species.</li> <li>Land-uses and densities must comply with NMBM Rural Management and Urban Edge Policy (e.g. 1 unit per 100 ha for infrastructure development, unless existing infrastructure already exists, individual footprints per unit to be limited to 1500 m²).</li> </ul>

	Pê	
SRK Consulting	Draft Bioregional Plan	

Low Density Rural Residential		MO.
Includes: development for 'lifestyle' or investment-type recreational ownership such as	Resort Zone	
share-block schemes, multi-ownership reserves, but only for extensive land portions with	WITH	This land-use type includes:
limited development (NB: excludes golf estates).	Nature Reserve	i. Various types of multi-owner or single-owner nature reserves;
Assumes the following conditions / controls:		ii. Share-block and sectional title developments in natural areas provided
<ul> <li>Maintenance of a large measure of natural land cover and ecological management</li> </ul>		the density of development is low (<20% development / degradation of
best practice implemented; densities of the development footprint should as per the		entire area); and
NMBM Rural Management and Urban Edge Policy;		iii. Properties and other extensive 'leisure and lifestyle' developments
<ul> <li>Individual footprints per unit as per the NMBM Rural Management and Urban Edge</li> </ul>		based on natural landscapes. Importantly, it excludes intensive
Policy (e.g. 1,500 m²);		commercial recreation-based developments such as resorts, eco-
<ul> <li>No additional rights should be granted if these will facilitate development outside of</li> </ul>		estates, and golf and polo estates. These latter approaches involve a
the parameters specified in points 1 & 2;		high proportion of habitat loss and very high investments in
<ul> <li>Clustered layout, sensitively placed to limit the development impact. Development</li> </ul>		infrastructure. They are often in effect urban development's outside the
within already transformed or disturbed areas or use of existing buildings or		urban edge and fall more appropriately into land-use type 9 (see
construction on timber piers (this will also ensure corridor linkages throughout the		below), which should be located within the urban edge or in peri-urban
property);		Zones.
<ul> <li>Sustainable water supply is guaranteed within the allocated Reserve of the water</li> </ul>		
resource;		
<ul> <li>Use off-grid services (solar power, rainwater harvesting, grey water recycling etc.)</li> </ul>		
unless services readily available nearby;		
<ul> <li>Architecture sympathetic to surrounding natural environment;</li> </ul>		
<ul> <li>Limited landscaping / gardening and only with locally indigenous plants;</li> </ul>		
<ul> <li>Rezone areas to Open Space 3 and declare as a Nature Reserve or Protected</li> </ul>		
Environment in terms of NEM: PA, and		
<ul> <li>Management agreement with NMBM or ECPB for conservation areas.</li> </ul>		

SRK Consulting Draft Bioregional Plan

# Table 4-4: Definitions of moderate impact rural land-use types

MODERATE IMPACT RURAL LAND-USE TYPES		
Land-use type	Example of zoning in terms of the new NMBM Zoning Scheme (not existing land-use rights)	Biodiversity Impact
Extensive Sustainable Livestock Production	Agriculture Zone	MODERATE
Includes: Mainly cattle, goat and sheep production on extensive areas of natural areas; mixed	AND	
livestock / wildlife options; tourism.	Resort Zone – where existing /	The ecological impacts of Extensive Livestock Production and Extensive
Assumes: Ecologically and economically sustainable management applied to farm portions above a certain minimum size, based on ecological and economic viability. Agricultural areas covered with natural vegetation, providing for sustainable low-impact agriculture-related landuses.		Game Farming can be similar and are dependent on the quality of veld management, i.e. a well managed cattle ranch may have less negative impact than a badly managed game farm, whereas with good management the reverse would be expected.
Stringent management conditions applied, such as -		of original representations of the production of
<ul> <li>Fire management measures, if necessary;</li> </ul>		nowever, because Extensive Livestock Production does not closely mining the previous nature state, which extensive dame farming does, it is recognized as
<ul> <li>Ecologically sustainable stocking rates (i.e. potentially lower than standard stocking rates);</li> <li>If mixed with wildlife, only indidenous or ecologically benion extra-limital species permitted:</li> </ul>		having a slightly higher impact. These areas can also include sustainable low-
<ul> <li>Resting cycles (rotational grazing);</li> </ul>		Inpact agricului e-Felateu Ianu-uses, e.g. mugenous piancha vesung.
<ul> <li>Avoidance of areas containing RDB species;</li> </ul>		
<ul> <li>Sensitive areas (e.g. wetland and river banks) to be protected from over-grazing and trampling, which can cause bank erosion and destabilisation, with subsequent degradation of water quality. The access areas for livestock should be demarcated and all alien invasive plants within this zone should also be eradicate; and</li> </ul>		
<ul> <li>Development and implementation of an Environmental Management Plan.</li> </ul>		
Densities and development footprints as per the NMBM Rural Management and Urban Edge Policy apply.		
Resort	Resort Zone	MODERATE
<u>Includes:</u> Holiday accommodation, Private Open Space for large-scale tourism purposes, nature conservation. The objective of the resort is to promote tourist and holiday facilities in areas with special environmental or recreational attributes, and to encourage access to these	WITH Open Space Zone 1 or 3	Resorts of this kind have a moderate impact due to the large scale nature of the development relative to Low Impact Eco-Tourism type developments. This land one and some of place to
tacilities by the general public.		ensure sustainability of the tourist attraction.
STEW/reeb Draft Bloreg	Draft Bioregional Plan 2 Dec 10 for gazetting (2014-final)	December 2010

att Bioregional Plan 2 Dec 10 for gazetting (2014-final)

Page 39

Small holdings  Includes: Agriculture, dwelling house, home and occupation. Such properties often occur close to cities, towns and villages, and may only occur within a demarcated urban edge or peri-urban zone. Development of this type should conform to the NMBM Rural Management and Urban Edge Policy for smallholdings near the urban edge. Some of these criteria prevent the further sub-division of small holdings in certain areas.  Small holdings of a residential character should ensure the maximum protection of indigenous veld, limited cultivation for household purposes, utilisation of existing buildings of development footprints where possible, use of existing infrastructure (roads) where possible and limited densification and expansion of the development footprint as per the NMBM Rural Management and Urban Edge Policy  Note - Applications in terms of Act 70 of 1970 for sub-division of residential type small holdings, which are zoned for agricultural purposes and located outside the urban edge, are subject to authorization from the Department of Agriculture to prevent the loss of agricultural land and fragmentation thereof. The NEMA EIA Regulations identify the subdivision of property of 9 ha or greater to 5 ha or less as a listed activity requiring a Basic Assessment.	Agriculture Zone or Residential  Zone 1 within:  Peri-urban Zone 1 & 2:  Smallholdings  Or  Rural Zone 1: Smallholdings  (RZ1)	Smallholdings do deplete natural resources, but can maintain areas of intact vegetation, amongst areas allocated for vegetables, horse paddocks or other farm animals. The small size of the property in the larger landscape does tend to result in fragmentation of natural areas, reducing the movement of wild animals and important pollinators. Smallholdings increase the extent of cumulative impacts as they require infrastructure (e.g. roads, pipelines & power lines) often beyond existing urban infrastructure.  Agricultural small holdings will often have a higher impact on biodiversity than rural residential small holdings, as the purpose will generally be for more extensive agricultural activities that are more destructive, but may still conserve the integrity of the natural environment and prevent the loss of agricultural land.

SRK Consulting

Draft Bioregional Plan

Table 4-5: Definitions of high impact agricultural land-uses

Land-use type	Example of zoning in terms of the new NMBM Zoning Scheme	Biodiversity Impact
HIGH IMPACT AGRICULTURAL LAND-USES	(not existing land-use rights)	
Arable Agriculture (forestry, dry land & irrigated cropping) with associated Rural	Agriculture Zone	noin
Settlement	within Rural Zone 2	Loui
Forestry (Timber Production) -		These farming activities have significant impacts on biodiversity. They
Includes: All timber plantations, mainly Pinus spp.		are mutually incompatible with biodiversity conservation, often accelerating
Assumes: Monoculture of alien timber species with heavy impact on hydrology and soil erosion		degradation by causing extensive habitat loss, soil erosion and hydrological changes. Their impacts vary from moderate to significant depletion of natural
and for introduction and spread of a variety of the most aggressive alien invasive plants.		flora & fauna and associated ecosystem functioning. Irrigated Crops and
Irrigated Crop Cultivation -		Timber Production impose particularly heavy impacts on environmental
Includes: All inicated crons (venetables) and inicated tree crons (crobards)		services such as water production. Their actual impact can be considerably
illolades. All Illigated Globs (Vegetables) alla Illigated tree Globs (Glorialus).		reduced by factors such as small scale, dispersed and sensitive placement
Assumes: Intensive production activity with high nutrient and agro-chemical inputs and often		and general good land husbandry.
two crops per year, ploughing, with no use of chemicals, results in irreversible loss of natural		Timber, certain other crops and alien plant infestations substantially lower
habitat.		water-tables and reduce stream flows. These land-uses consume large
Dryland Crop Cultivation -		quantities of water potentially available for other uses, including the minimal
		flow required for ecosystem maintenance – i.e. the 'ecological reserve'. As a
Includes: All tillage cultivation of non-irrigated crops, mostly single-season annuals, but		result, a complex authorisation protocol is in place to issue water-use
including perennial and orchard-type tree crops if cultivated with an indigenous grass layer.		licenses to approved land users wanting to grow such crops (see 'water
Assumes: Crop production methods that conserve water and protect against soil erosion;		extraction' above). The licensing process is based on EIA procedures.
limited and responsible use of fertilisers, pesticides and other agrochemicals and Genetically		In Contrast -
Modified Organisms (GMOs).		These land-use types can be improved by maintaining natural corridors
		within the landscape and adopting biodiversity-friendly agricultural / farming
		practices e.g. water conservation, organic fertilizers, minimal tillage etc.
		Irrigated and dryland crop cultivation can provide opportunities for movement of species and connectivity between CBAs under certain conditions.
Intensive Animal Farming with associated Rural Settlement	Agriculture Zone	
STEWireb Draft Boregi	Draft Bioregional Plan 2 Dec 10 for gazetting (2014-final)	December 2010

Page 41

Impact of <b>dairy farming</b> and similar production systems can be reduced by not developing planted pastures and applying:	<ul> <li>Ecologically sustainable stocking rates (i.e. potentially lower than standard stocking rates);</li> <li>Resting cycles (rotational grazing);</li> <li>Avoidance of areas containing Red Database species;</li> <li>Sensitive areas (e.g. wetland and river banks) to be protected from overgrazing and trampling, which can cause bank erosion and destabilisation, with subsequent degradation of water quality. The access areas for livestock should be demarcated and all alien invasive plants within this zone should also be eradicate; and</li> <li>Development and implementation of an Environmental Management Plan.</li> </ul>
with consent use required within Rural Zone 2	
Includes: All intensive animal production systems, of domestic or 'wild' species that are primarily dependent on imported foodstuffs and confinement; dairy farming and all areas in production support for dairy, including pastures, fodder and grain crops, much of which is usually irrigated.	Assumes: Intensive production areas with relatively dense development; not dependent on the use of natural vegetation for production and associated settlements for the farming work force.

an 2 Dec 10 for gazetting (2014-final)

qe

SRK Consulting Draft Bioregional Plan Table 4-6: Definitions of high impact urban and industrial land-uses

Land-use type	Example of zoning in terms of the new NMBM Zoning Scheme (not existing land-use rights)	Biodiversity impact
MIGH IMPACT URBAN & INDUSTRIAL LAND-USES		
Residential, Business & Industrial i.e. Urban land-uses	Residential Zone 1, 2, 3 or 4	VERY HIGH
<u>Includes:</u> All residential, commercial, industrial and tourism development at densities typical of urban or peri-urban areas.	Business Zone 1, 2, 3 or 4 Industrial Zone 1 or 2	These land-uses destroy and / or sterilise natural vegetation and soil. In urban areas whole landscapes are modified in this way. Where biodiversity
Assumes: Peri-urban development, termed 'rural residential', 'eco-estates', or 'lifestyle developments' by developers; includes resorts, & golf estates.	Community Zone 1 or 2	exists or is deliberately protected, such as in urban nature reserves, it can be negatively affected by its small scale and ecological isolation.
	Transport Zone 1, 2 or 3 Undetermined	Dams built for water supply can have significant impacts on aquatic biodiversity by changing hydrological regimes, such as water volume, velocity and temperature, which modify downstream habitats e.g. freshwater
Infrastructure (roads, power lines, pipelines)	Transport Zone 1, 2 or 3	inflow into estuaries required for organisms present in the estuary, natural
Includes: All hardened roads, railways, canals and pipelines, including power and telephone lines, which can have a measurable impact on habitats and particular species e.g. power lines on birds.		rivers creates barriers to fish migration routes. Off channel construction may also impact negatively on important terrestrial biodiversity.
Mining & Quarrying  Includes: All strip and opencast mining excavations or quarrying (sand mining); plus the visual, physical and chemical impacts of these activities, particularly on ground water reserves; all mine waste and refuse dumps, urban waste sites and landfill sites for whatever purposes.	Industry Zone 3: Extractive Industry	In Contrast:  Metropolitan Open Space Systems – can improve the conservation of biodiversity in urban & peri-urban areas by maintaining natural corridors, thereby creating a biodiversity network for important ecosystem services.
<u>Dams</u> <u>Includes:</u> All substantial impoundments, dams and weirs, with associated pipelines, canals, access roads and bulk water transfer schemes.	Agricultural Zone	Mining should be undertaken in a phased manner to allow restoration or rehabilitation of disturbed areas over the length of the project.  Infrastructure e.g. pipelines, roads etc. should be designed to avoid Critical Biodiversity Areas and aligned with existing infrastructure or on developed / degraded footprints where nossible

# 5 Part E: Other measures for effective management

#### 5.1 Planning

#### 5.1.1 Protected Area Expansion

Only about 2.4 % (approximately 4,700 ha) of the municipal area enjoys some form of statutory protection. Three additional formal protected areas are proposed, namely the van der Kempskloof (778 ha), Chatty (1,161 ha), and Aloes (226 ha) Nature Reserves. However, the inclusion of these areas will not be a sufficient to secure a representative proportion of the municipality's biodiversity.

The NMBM owns a relatively limited amount of land (approx. 72,700 ha) within the municipal area, with the majority of its 195,400 ha being under private ownership (62.78 %). Since insufficient funds exist for NMBM to acquire and manage all of the property required for conservation, protected area expansion will require the conservation of a combination of privately-owned and municipally-owned land parcels. A range of mechanisms need to be developed in order to support protected area expansion within the municipal area, including the implementation of biodiversity offsets and the development of conservation agreements with private landowners. The rezoning of municipally owned properties located within the network of Critical Biodiversity Areas and Ecosystem Support Areas to Open Space III is a further priority.

#### **Biodiversity Offsets**

Where development proposals will result in biodiversity impacts after mitigation that range from moderate to severe, biodiversity offsets should be considered to compensate for the residual impacts of the development. To date, biodiversity offsets have only been used in a few cases in the municipality, but due to the extent of habitat loss during the last five years, the need to consider the application of biodiversity offsets is expected to increase substantially.

Biodiversity offsets, where appropriate, can comprise either single or composite areas in order to compensate adequately for residual biodiversity loss. An acceptable measure of the residual loss is obtained as a point of departure for determining an appropriate biodiversity offset. Biodiversity offsets are calculated by multiplying this measure by a basic biodiversity offset ratio linked to the Ecosystem Threat Status of the affected ecosystem.

The following ratios are recommended with regard for biodiversity offsets (DEA&DP, 2008):

- A 30:1 ratio for *Critically Endangered* ecosystems and *Critical Ecosystem Process Areas*, where a biodiversity offset would be appropriate in exceptional circumstances only;
- A 20:1 ratio for Endangered ecosystems;
- A 10:1 ratio for Vulnerable ecosystems; and
- No biodiversity offset for *Least Threatened* ecosystems.

The area determined by the basic biodiversity offset ratio should then be adjusted by a range of context-specific considerations, such as:

STEWreet

Draft Bioregional Plan 2 Dec 10 for gazetting (2014-final)

Page 44

- The condition of the affected habitat;
- The significance of residual impacts on threatened species;
- The significance of residual impact on special habitats;
- The significance of residual impact on important ecological corridors or process areas; and
- The significance of residual impact on biodiversity underpinning valued ecosystem services.

For a biodiversity offset to contribute effectively to protected area expansion and biodiversity conservation in the Nelson Mandela Bay Municipality, biodiversity offsets should be located in a 'biodiversity offset receiving area', which in the context of the NMBM will be the Critical Biodiversity Areas.

#### **Biodiversity Stewardship**

The NMBM Environmental Services Business Unit is currently developing a programme based on the biodiversity stewardship approach currently implemented in the Western Cape and KwaZulu-Natal, and adopted by the National Biodiversity Framework (2009) as a preferred mechanism for protected area expansion. The programme will provide for formal, legally-recognised agreements (potentially also declaration of land as Nature Reserve and Protected Environments in terms of the Protected Areas Act) between the municipality and landowners to conserve biodiversity. In order to support this programme, the NMBM has adopted a property rates policy that will allow for rebates for private landowners who enter into agreements with the municipality to conserve their land.

Current private landowner conservation initiatives are primarily centred in the Sardinia Bay, Bushy Park area (i.e. Sardinia Bay Conservancy) and Blue Horizon Bay areas. However, in most instances legal protection of natural habitats of high conservation value outside of statutory protected areas in Nelson Mandela Bay is not in place.

A formal protected area expansion strategy has not been developed for the Nelson Mandela Bay and the NMBM currently has no targets set for protected area expansion. Spatial priority areas for protected areas expansion should be aligned to the network of Critical Biodiversity Areas described in the bioregional plan.

The following recommendations are made with regard to protected area expansion:

- A protected area expansion strategy should be developed for the province and refined and adapted for the municipality;
- The development of the NMBM's biodiversity stewardship programme should be supported as
  this approach is recognised as being one of the most cost-effective mechanisms for protected areas
  expansion;
- In order to improve the status of the local authority reserves in the municipal area, they should be assessed in terms of their biodiversity value and the requirements of the Protected Areas Act and then declared as either Nature Reserves or Protected Environments based on this assessment;
- The corridor between the Maitlands Nature Reserve and the Island Nature Reserve should be consolidated;

Page 45

- The coastal corridor between Sardinia Bay Nature Reserve and Nelson Mandela Metropolitan
  University Private Nature Reserve should be consolidated. The NMMU should be encouraged to
  seek declaration for their reserve in terms of section 23 of Protected Areas Act;
- The entire extent of municipal land within the Baakens Valley should be declared as a Nature Reserve in terms of Protected Areas Act. Private owners of land along the valley should be encouraged to enter into biodiversity stewardship agreements with the NMBM;
- The corridor between the Van Stadens Wildflower Reserve, Lady Slipper Nature Reserve and Van Stadensberg Natural Heritage Site should be consolidated. The Van Stadensberg Natural Heritage Site should be declared as a Nature Reserve in terms of the Protected Areas Act. The Van Stadens River Conservancy initiative should be supported by the NMBM biodiversity stewardship programme in an effort to link the Van Stadens Wildflower Reserve to the coast;
- Protected Environment status should be investigated for the Sardinia Bay, Bushy Park and Horizon
  Bay area's to support the efforts of landowners and to restrict certain activities which would be
  detrimental to the important biodiversity of these areas;
- The NMBM land at Van Der Kemp's Kloof / Parsonsvlei / Chatty Kloof should be declared as a Nature Reserve. The Anglican Church land should be incorporated into this Nature Reserve through a biodiversity stewardship agreement. This Nature Reserve should link to the Joe Slovo Thicket Corridor and the proposed Hopewell Nature Reserve, which in turn should be linked to the Groendal Wilderness area; and
- The corridor between the Springs Nature Reserve and the Groendal Wilderness area should be consolidated.

#### 5.1.2 Ecosystems and Species Requiring Biodiversity Management Plans

Biodiversity management plans are one of the planning tools provided for by the Biodiversity Act. The act provides for any person, organisation or organ of state to submit to the Minister a draft biodiversity management plan for an ecosystem or indigenous species which warrants special conservation species. Before approving a draft biodiversity management plan, the Minister must identify a suitable person, organisation or organ or state which is willing to be responsible for the implementation of the plan. Importantly, the existence of a biodiversity management plan for an ecosystem or species enables the Minister to enter into a biodiversity management agreement with a person, organisation or organ regarding its implementation. Landowners that are party to a biodiversity management agreement are eligible for the income tax deductions, as provided for by the Income Tax Act (Act 58 of 1962), as amended by the Revenue Laws Amendment Act (Act 60 of 2008). The development of biodiversity management plans in terms of Biodiversity Act is therefore a useful pursuit to support the NMBM's emergent biodiversity stewardship programme.

Within the NMBM area the following ecosystems are recommended for the development of biodiversity management plans in terms of the Biodiversity Act:

• The Swartkops River system (Implementation site 13) – due to the value of this system in terms of both the ecological processes that occur and the environmental goods and services that it provides, and due to the threats this system faces; and

STEWreeb

Page 46

• Urban lowland fynbos (Implementation site 2) – including the *Critically Endangered* Algoa grassy fynbos complex ecosystems: Walmer Grassy Fynbos, Baakens Grassy Fynbos, Lorraine Transitional Grassy Fynbos and Colleen Glen Grassy Fynbos systems. The natural extent of these ecosystems has been severely reduced by urban development and, without appropriate management intervention, these systems may collapse.

Species that are recommended for the development of biodiversity management plans in terms of the Biodiversity Act are:

- Cyclopia pubescens (Critically Endangered) a rare and highly localised species, occurring from
  the foot of the Van Stadens Mountains to just west of Port Elizabeth, which is under threat from
  urban expansion. A management plan for this species was produced by the NMMU Centre for
  African Conservation Ecology;
- Aloe bowiea (Critically Endangered) a highly restricted species, only known from three sites
  near Motherwell, that is threatened by expansion of informal and formal housing, harvesting,
  overgrazing;
- Aspalathus recurvispina (Critically Endangered) thought to be extinct. The species rediscovered
  in the wild near Humewood, but is highly threatened by habitat loss through urban development.
  Only a single population of this plant is known to occur;
- Orthopterum coegana (Critically Endangered) a succulent that was restricted to three
  populations occurring on sandstone outcrops around Coega Kop, east of Port Elizabeth. One of
  these populations had to be relocated when the western portion of Coega Kop was targeted for
  mining for the construction of Coega deepwater port;
- Fitzsimons whip lizard Tetradactylus fitzsimonsi is restricted to the Algoa basin proposed as a
  Critically Endangered species; and
- Albany adder Bitis albanica one of South Africa's rarest snakes. It is restricted to the Algoa
  basin, threatened by habitat loss, principally due to mining, and has been proposed as Critically
  Endangered.

Biodiversity management plans should be developed in conjunction with the relevant NMBM Conservation Area Managers, and actions should be incorporated into the key performance indicators for managers and into Environmental Management's Service Delivery Improvement Plan.

### 5.2 Operation and Management

There is a need to increase budgetary provision to enable the NMBM to adequately meet its ecological management obligations.

Management actions that are required to effectively conserve biodiversity within the municipal area include invasive alien species management and rehabilitation and restoration of impacted areas.

#### 5.2.1 Invasive Alien Species Management

An invasive alien plant clearing strategy was developed for the NMBM in April 2009 (SSI, 2009). Areas that were identified as priorities were the Swartkops River, Chatty River, Baakens River, Paapenkuils River, Sardinia Bay and Vans Stadensberg corridor – all Critical Biodiversity Areas. The

STEWreeb

Draft Bioregional Plan 2 Dec 10 for gazetting (2014-final)

Page 47

strategy schedules areas within the priority areas for clearing, and provides an estimate of the cost and effort to clear.

The following is recommended with regard to invasive alien species management:

- The NMBM should ensure that adequate budget is provided for invasive alien plant control
  operations on municipal land;
- Private landowners should be encouraged to control invasive alien plants on their properties (e.g.
  through the provision of property rates rebates for landowners who adequately control invasive
  alien plants on their properties); and
- The NMBM's invasive alien plant clearing strategy, which is aligned to the bioregional plan, should be used to direct investment in alien plant control.

#### 5.2.2 Priority Areas for Rehabilitation and Restoration

The following areas are recommended as priorities for rehabilitation and restoration in order for these areas to support the network of critical biodiversity sites within the NMBM municipal area:

- The Swartkops River system areas along the KwaZunga River (upper reaches of the Swartkops River system) have been impacted by sand-mining, which has been described as the second-most significant impact on the river and has downstream impacts on the Swartkops River (IWR, undated);
- The Parsonsvlei / van der Kempskloof / Chatty Kloof / Hopewell area this area has been impacted by mining and the old and existing quarries will require rehabilitation. Dumping of domestic rubble is particularly bad in this area and efforts should be made to clear this and prevent its reoccurrence;
- The coastal belt, particularly from the Sardinia Bay and Sylvic Nature Reserves to the Cape Receive and NMMU Nature Reserves is highly invaded by invasive alien plants. Active rehabilitation may be required in this area on removal of the invasive alien plants in order to prevent accelerated soil erosion and destabilisation of dunes; and
- The Grassridge area is currently mined for limestone. This area houses Grassridge Bontveld and the Albany adder *Bitis albanica* and will require rehabilitation on completion of mining activities.

#### 5.3 Mainstreaming

#### 5.3.1 Institutional Arrangements

The NMBM has instituted an internal organisational structure, the Corporate EIA Task Team (CETT), to facilitate the coordination of EIA applications and broader environmental issues. CETT enables review by the various directorates within the NMBM of the findings of EIAs and allows for mainstreaming of environmental issues, including the NMBM Conservation Plan, into operations of the directorates. This existing structure can be used to mainstream biodiversity management considerations into the operations of the various municipal directorates for new projects.

A Multi-Disciplinary Technical Task Team (MDTTT) also exists to facilitate co-operative decision-making between the relevant municipal directorates on strategic projects. It is recommended that the relevant sub-directorates (e.g. Transportation Planning, Water and Sanitation etc.) be requested to

STEWreeb

Draft Bioregional Plan 2 Dec 10 for gazetting (2014-final)

Page 48

provide quarterly updates at the MDTTT meetings on proposed short, medium and long-term projects that may be in conflict with the bioregional plan or result in impacts on the network of Critical Biodiversity Areas. The MDTTT is also an ideal structure to identify opportunities for collaboration between directorates regarding socio-economic projects linked to biodiversity, and construction projects where rehabilitation and / or restoration are required.

Monthly bilateral meetings between the NMBM and DEDEA also take place and serve as a mechanism for coordination of EIA applications and broader environmental issues. These meetings provide further opportunity to address potential conflicts between proposed land-uses and the bioregional plan.

#### 5.4 Building a Biodiversity Economy

Building a biodiversity economy involves promoting economic activity that uses biodiversity in a sustainable way. This includes the creation of opportunities through biodiversity for economic development in previously disadvantaged communities, the promotion of sustainable agriculture and fisheries, and the promotion of land-uses that are compatible with biodiversity (for example, sensitive ecotourism and game farming).

#### 5.4.1 Ecotourism

The NMBM is unique in South Africa in that it is an area of convergence for five of South Africa's biomes. Thus, by visiting the NMBM, the eco-tourist can experience a small taste of what the country in a whole has to offer.

The NMBM has undergone a process of developing Conservation Development Frameworks (CDFs) for many of its local authority reserves. One of the functions of the CDFs is to identify areas within the reserves that are appropriate for sensitive development, such as ecotourism facilities. The NMBM will look to enter into partnerships with the private sector in order to unlock the economic potential of these reserves. Conservation Development Frameworks have been developed for the proposed Van Der Kemps Kloof / Chatty Nature Reserve, Aloes / Swartkops Nature Reserve, Settlers Park Nature Reserve, the Target Kloof / William Moffet section of the Baakens Valley, Maitlands Nature Reserve and the Van Stadens Wildflower Reserve.

The following is recommended with regard to the development of ecotourism within the municipal area:

- The concept of the "Five Biome City" should be promoted, potentially through the development of a route that exposes tourists to the various biomes within the municipal area; and
- The NMBM should implement mechanisms, such as the development of public-private partnerships, to unlock the ecotourism potential of its local authority reserves, specifically through the sensitive and appropriate development (e.g. restaurants, tea-gardens and lodges) of areas identified in the Conservation Development Frameworks.

#### 5.4.2 Nature-based Development Models

Recent years have marked a growth in the number of rural lifestyle residential developments within the municipal area. These developments have the potential to support biodiversity conservation objectives if they are appropriately planned and managed.

STEWreeb

Page 49

Development models that support the conservation of Critical Biodiversity Areas, are aligned with the NMBM's land-use policies, and are underpinned a by legitimate conservation management ethic should be encouraged.

#### 5.4.3 Game Reserves and Game Farming

The northern portion of the NMBM is well suited to the development of the game reserve and game farming industry. Farming with game in the thicket biome has the potential to be more ecologically and economically sustainable than farming with livestock. The main activity for which game is utilised is hunting, including both recreational and trophy hunting. With the Eastern Cape being recognised as the most preferred destination for overseas hunters (Smith & Wilson, 2002), this industry has potential to contribute to a biodiversity based economy in the NMBM, provided that it is correctly managed and controlled (for example, preventing or limiting the introduction of non-indigenous species).

The following is recommended with regard to game reserves and game farming:

Game reserves and game farming should be promoted in areas that are suitable for these activities
(areas within the thicket biome) within the NMBM, provided that such operations are managed in
a way that supports biodiversity conservation (e.g. stock only indigenous species and appropriate
densities).

#### 5.4.4 Skills Development and Training

The development of a biodiversity economy within the municipal area will create a demand for a new set of skills within the workforce. Skills development and training opportunities linked to biodiversity include those associated with the development of indigenous plant nurseries, ecotourism establishments and secondary industries centred on invasive alien plant biomass.

An opportunity for skills development is the creation of indigenous plant nurseries to supply plants for cultural and medicinal use. The nurseries should be linked to the establishment of a training college, potentially supported by the SANBI Green Futures Programme, in order to train previously disadvantaged individuals in horticulture. An additional benefit of the nurseries would be the reduced demand on natural systems for plants of cultural, landscaping and medicinal use.

Ecotourism establishments, such as game lodges, require staff skilled in the hospitality sector and in ecological management. Skills development programmes linked to biodiversity conservation that seek to train and employ youth displaced by HIV / AIDS and poverty should be encouraged.

The development of industries within the municipal area that utilise invasive alien plant material will result in some economic return for the investment spent on control operations and also the development of skills and creation of jobs. Opportunities exist to create new industries and produce various products from invasive alien plant biomass. The following is recommended with regard to skills development and training:

 Skills development models that link biodiversity conservation to improving the capacity of disadvantaged individuals should be encouraged.

#### 5.4.5 Natural Resource Use

A proportion of the NMBM's population makes use of natural resources for cultural and medicinal purposes, as a source of fuel, and as a means to supplement income. Unsustainable use of natural

STEWreeb

Draft Bioregional Plan 2 Dec 10 for gazetting (2014-final)

Page 50

resources will impact on biodiversity and on the ability of the environment to supply goods and services to the population.

The following is recommended with regard to natural resource use:

- Areas of natural vegetation, outside of the existing protected area network, should be set aside for traditional practises to continue;
- In order to reduce the impact of resource use on natural areas, the development of nurseries for indigenous plants of cultural and medicinal value should be encouraged; and
- The impacts of the bait collecting industry at Swartkops should be monitored and the industry should continue to be regulated to ensure that harvesting is sustainable.

#### 5.4.6 Payment for Ecosystem Services

Payment for Ecosystem Services is the practice of offering incentives to farmers or landowners in exchange for managing their land to provide some sort of ecological service. Amongst the ecosystem services receiving the most attention worldwide is climate change mitigation and, related to this, there is an opportunity to develop the carbon sequestration industry within the NMBM.

Spekboom *Portulacaria afra*, which is a dominant species in many thicket types, is being increasingly used to restore plant cover in degraded thicket. This plant has the ability to sprout from replanted cuttings, making it a good candidate for large-scale restoration of degraded land. Furthermore, there is evidence that spekboom has extraordinary carbon storing capabilities. Restoration of degraded subtropical thicket would achieve the combined aims of improving rural livelihoods, restoring biodiversity, and replenishing natural capital and ecosystem services (Powell *et al.*, undated). Already, private companies have begun offering services to support the restoration of thicket by landowners.

Opportunities for thicket restoration projects are in the north of the municipal area, in places where subtropical thicket has been degraded or transformed by human activities.

The following is recommended with regard to payment for ecosystem services:

• The development of thicket restoration projects within the NMBM should be encouraged.

#### 5.4.7 Expanded Public Works Programme

The Expanded Public Works Programme (EPWP) is a government programme to provide poverty and income relief through temporary work for the unemployed to carry out socially useful activities. Programmes active within the municipal area include the Working for Water, the Working for Woodlands and CoastCare Programmes.

The Working for Water Programme collaborates with local communities on projects to control invasive alien plants. At the same time, the programme creates an enabling environment for skills training and invests in the development of the communities through the implementation of HIV and AIDS projects and other socio-development initiatives. Short-term contract jobs (with an emphasis on recruiting women, youth, and the disabled) are created through the clearing activities undertaken.

The following is recommended with regard to Expanded Public Works Programmes within the NMBM:

Page 51

- The efforts of the EPWP programmes should, where possible be aligned to the network of Critical Biodiversity Areas.
- Where private landowners enter into conservation agreements with the NMBM, these landowners should be prioritised for assistance by EPWP programmes.

# 6 Part F: Monitoring, Reviewing and Updating

The Nelson Mandela Bay Municipality is the lead implementing agent of the bioregional plan for the bioregion. As such, the NBMM is responsible for overseeing the implementation of the plan and future review thereof, including:

- Its application in land-use planning and decision-making;
- Revision and amendment of the underlying systematic biodiversity plan (Conservation Assessment and Plan for the Nelson Mandela Bay Municipality) at least every five years; and
- An update of the published bioregional plan at least every five years.

However, it is important to note that SANParks, the Eastern Cape Parks, and private landowners (e.g. Mountains to Oceans, PPC, Coega Development Corporation etc.) are responsible for the management of certain protected areas within Nelson Mandela Bay and the above lead agency responsibilities assigned to the NMBM do not in any way remove or transfer existing and future management responsibilities from these management agencies.

#### 6.1 Review

Prior to the commencement of a full revision of the bioregional plan that is in force at the time, it is recommended that the plan be reviewed in order to identify key components that require updating.

Furthermore, the NMBM has put various organisational structures in place to facilitate the integration of the NMBM CBA Network into land-use planning and decision-making (see Section 5.3.1). The effectiveness of these structures in achieving this objective should be reviewed bi-annually. The land-use guidelines (see Section 4) associated with this bioregional plan are additional instruments to support the application of the plan in planning and decision-making. The guidelines should be reviewed at the same intervals as the bioregional plan.

#### 6.2 Monitoring

Various monitoring instruments and indicators were considered, in addition to the availability of necessary resources to measure achievement of the *Planning*, *Operation and Management*, *Mainstreaming* and *Building a Biodiversity Economy* objectives of the bioregional plan. It is recommended that monitoring be undertaken on a five-yearly cycle, in alignment with the revision of the bioregional plan.

Implementation of the various recommended measures for effective management and monitoring should commence immediately upon gazetting of the bioregional plan.

STEWreeb

Page 52

#### 6.2.1 Monitoring Instruments

The following instruments were identified as being viable to generate reliable data for monitoring purposes:

- Previous and future systematic biodiversity plans;
- NMBM Alien Invasive Plant Control Strategy;
- NMBM Environmental Management Sub-Directorate SDIP;
- Biodiversity Management Plans;
- Protected Area Management Plans;
- NMBM State of the Environment Reports;
- Biodiversity Stewardship agreements;
- Development of a system of tracking NEMA EIA Regulations and MPRDA applications that do
  not correctly refer to the NMBM bioregional plan, as well as NMBM project proposals that present
  conflicts with the Bioregional Plan that have not been evaluated via the MDTTT;
- Future NMBM Protected Area Expansion Strategy; and
- Progress Reports for NMBM and other biodiversity projects in the municipal area.

#### 6.2.2 Monitoring Indicators

Due to the limited number of instruments available to generate reliable data for monitoring purposes, only the following final indicators were selected per category (DEAT, 2005; SANBI, 2009):

#### **Planning**

- Protection status of ecosystems (hectares per vegetation type and protected area categories);
- Terrestrial Habitat Loss (hectares / vegetation type);
- Threatened amphibian species (hectares and number of populations / threatened species);
- Threatened bird species (hectares and number of populations / threatened species);
- Threatened mammal species (hectares and number of populations / threatened species);
- Threatened reptile species (hectares and number of populations / threatened species);
- Biodiversity management plans for ecosystems (number of plans);
- Biodiversity management plans for species (number of plans);
- · Biodiversity Stewardship Framework developed; and
- Protected Area Expansion Strategy developed.

#### **Operation and Management**

- Budgetary provision for conservation management (percentage in relation to required budget);
- Area of invasive alien species cleared (hectares / Ecosystem Threat Status or CBA-ESA category);
   and

STEWreeb

Page 53

 Area of natural vegetation restored or rehabilitated (hectares / Ecosystem Threat Status or CBA-ESA category).

#### Mainstreaming

- EIA and MPRDA applications that do not refer to the NMBM bioregional plan appropriately (percentage); and
- Municipal project proposals that are in conflict with the bioregional plan, which have not been reviewed by the MDTTT (percentage).

#### **Building a Biodiversity Economy**

- Temporary and permanent employment opportunities created through biodiversity initiatives (number of jobs);
- Protection status of ecosystems and species as a direct result of development projects (hectares / vegetation type or species);
- Individuals trained in biodiversity-related industries (number); and
- Areas restored or rehabilitated as a direct result of payment for ecosystem services (hectares / vegetation type or species).

#### 6.3 Updating

The updating of a systematic biodiversity plan is a data intensive and time consuming process. In light of the requirement for municipal SDFs to undergo full revisions every five years, it is recommended that the NMBM systematic biodiversity plan also be updated on a five-yearly cycle. The revision should be scheduled such that a draft (revised) biodiversity plan is completed by the commencement date of the SDF revision process; thereby facilitating an integrative planning process and the incorporation of the Critical Biodiversity Areas map into the SDF.

Due to ongoing changes in land-use rights and the implications of development on the NMBM CBA network, a biennial update of the Critical Biodiversity Areas map should also be considered. The revision should be made available as a digital map and in GIS via the NMBM's website. The revised map must be submitted to the MEC of DEDEA, in consultation with the SANBI and the Minister of DEA, for gazetting.

Updates of the entire bioregional plan should be undertaken on a five-yearly cycle, once the revision of the NMBM systematic biodiversity plan has been completed. However, more frequent updates to specific components (other than the CBA map) could be undertaken when necessary.

# 7 Part G: GIS Files

A data package of the principal spatial layers that were used in the development of the bioregional plan has been assembled. For ease of reference, an outline of the shapefiles (.shp), associated layer files (.lyr), and important fields in the attribute tables is included in Table 1. The shapefiles were created in ESRI ArcMap 9.3.1 and include metadata (.xml files) according to the ISO Metadata Standard, which conforms to the SANBI Biodiversity GIS metadata standards.

#### Table 7-1: Outline of the shapefiles included in the data package

STEWreeb

Draft Bioregional Plan 2 Dec 10 for gazetting (2014-final)

Page 54

Folder	Shapefiles	Layer files	Description	Important fields	Description
FINAL_CRITICAL_ BIODIVERSITY AREAS_2010	400919_NMBM_CRITIC AL_BIODIVERSITY_AR EAS_FINAL_07122010. shp	400919_NMBM_CR ITICAL_BIODIVER STIY_AREAS_FINA L_07122010.lyr	Final shapefile of Critical Biodiversity Areas (CBAs) for the NMBM	CONSERVATN	Classification of the CBAs
PROTECTED_ARE AS_2009	PROTECTED_AREAS_ 2007.shp	367380_NMBM_PR OTECTED_AREAS _161107.lyr	The statutory protected areas and non-statutory conservation areas of the NMBM	RESNAME	Protected area name
ECOSYSTEM_STA TUS_2009	367380_VEGTYPES_FI NAL_050309.shp	367380_NMBM_VE GETATION_TYPES _2009.lyr	The vegetation types of the NMBM	VEGTYPE	The vegetation type classification
		367380_NMBM_EC OSYSTEM_STATU S_2009.lyr	The Ecosystem Threat Status classification of the NMBM vegetation types	ECOSYSTEM_	The Ecosystem Threat Status classification
SPECIES_2009	367380_SPECIES_OF_ SPECIAL_CONCERN_ NUMBERED_010307.sh p	-	The species of special concern layer for the NMBM. In order to prevent misuse of the data, the names of the species are not shown – a code is instead assigned to each species of special concern	SPP_NUMBER	Reference code for the species of special concern occurring
CRITICAL_ECOLO GICAL_PROCESS ES_2009	367380_NMBM_CRITIC AL_ECOLOGICAL_PRO CESSES_051007.shp	367380_NMBM_CR ITICAL_ECOLOGIC AL_PROCESSES_ 161107.lyr	The critical ecological processes within the NMBM	ECOSYSTEM_	Ecological process type classification

# 8 Conclusions and Recommendations

The draft bioregional plan has identified priority areas for biodiversity conservation in the Nelson Mandela Bay municipal area. Measures for effective management, review, monitoring and updating have also been recommended. Implementation of the various recommended measures for effective management and monitoring should commence immediately upon gazetting of the bioregional plan.

The bioregional plan serves as the official reference for biodiversity priorities within the bioregion, superseding all the previous non-published spatial biodiversity plans for the area. Upon gazetting of the bioregional plan, all organs of state will be required to take the bioregional plan into account in their decision-making.

This draft bioregional plan is consistent with the Biodiversity Act and meets all the requirements of the *Guideline regarding the Determination of Bioregions and the Preparation and Publication of Bioregional Plans* (DEAT, 2008) for review by a Ministerial Panel.

Page 55

SRK Consulting Catifold Floreness Synthetics

SRK Consulting

Body of Tablepox

8276-9694-3083-STEW
This signature has been privated digitally. The Author has given permises use for this document The details are stored in the SRX Signature Data

Warrick Stewart M.Sc. Pr.Sci.Nat.

Senior Environmental Scientist

Brian Reeves M.Sc. Pr.Sci.Nat.

Senior Environmental Scientist

CONSTITUTED

AUTOMOTIVE TO THE CONTROL OF THE CONTR

Rob Gardiner M.Sc. Pr.Sci.Nat.

Principal Environmental Scientist

**SRK Consulting** 

Page 56

## 9 References

ADU. 2009. Seabird sites in southern Africa: Cape Recife, Mandela Bay, Port Elizabeth. Avian Demography Unit website: <a href="http://adu.org.za/cprecife.php">http://adu.org.za/cprecife.php</a>

Berliner, D. & Desmet, P. 2007. Eastern Cape Biodiversity Conservation Plan: Technical Report. Department of Water Affairs and Forestry Project No 2005-012, Pretoria. 1 August 2007.

Bornman, T.G. & Klages, N.T.W. 2003. Maitland River Mouth Strategic Environmental Assessment. Institute for Environmental and Coastal Management. IECM Report No. C81, University of Port Elizabeth, Port Elizabeth.

Bruton, N. 2006. Nelson Mandela Metropole Business Confidence Indicator: October 2006. Port Elizabeth Regional Chamber of Commerce and Industry, Port Elizabeth.

ACE. 2008. Centre for African Conservation Ecology: Annual Report 2008. Centre for African Conservation Ecology, Nelson Mandela Metropolitan University, Port Elizabeth.

CI. 2009. Biodiversity Hotspots. Conservation International website: <a href="http://www.biodiversityhotspots.org/Pages/default.aspx">http://www.biodiversityhotspots.org/Pages/default.aspx</a>

Cowling R.M., Pressey R.L., Lombard A.T., Desmet P.G. & Ellis A. 1999a. From representation to persistence: requirements for a sustainable system of conservation areas in the species-rich mediterranean-climate desert of southern Africa. Diversity and Distributions 5, 1-21.

Cowling R.M., Pressey R.L., Lombard A.T., Heijnis C.J., Richardson D.M. & Cole, N. 1999b. Framework for a conservation plan for the Cape Floristic Region. IPC Report 9902 submitted to WWF:SA.

Cowling, R.M., Lombard, A.T., Rouget, M., Kerley, G.I.H., Wolf, T., Sims-Castley, R., Knight, A., Vlok J.H.J., Pierce, S.M., Boshoff, A.F. & Wilson, S.L. 2003. *A Conservation Assessment for the Subtropical Thicket Biome*. Report 43 to the Terrestrial Ecology Research Unit, University of Port Elizabeth, South Africa.

CSIR. 2004. Eastern Cape State of the Environment Report. CSIR Division of Water, Environment and Forestry Technology. Durban, South Africa.

DEAT. 2005. South Africa's National Biodiversity Strategy and Action Plan. Department of Environmental Affairs & Tourism, South Africa.

Department of Environmental Affairs and Development Planning. 2007. *Provincial Guideline on Biodiversity Offsets*. Republic of South Africa, Provincial Government of the Western Cape, Department of Environmental Affairs & Development Planning, Cape Town.

IWR. Undated. Catchment M: The Swartkops River. Institute for Water Research, Rhodes University, Grahamstown.

Jenkens, A.R. & Stephenson, A. Siberian Peregrines in southern Africa. *Africa – Birds & Birding* 2(4): 63-68.

Lubke, R.A. & de Moor, I. 1998. A Field Guide to the Eastern & Southern Cape Coasts. University of Cape Town Press, Cape Town.

Page 57

Martin, P. 2002. South African Birding: Birding Resources for Southern Africa. SA Birding website: <a href="http://www.sabirding.co.za/birdspot/021017.asp">http://www.sabirding.co.za/birdspot/021017.asp</a>

Mittermeier, R. A., Robles Gil, P., Hoffmann, M., Pilgrim, J., Brooks, T., Goettsch Mittermeier, C., Lamoreux, J. & Da Fonseca, G.A.B. 2004. Hotspots Revisited: Earth's Biologically Richest and Most Endangered Terrestrial Ecoregions. Conservation International, Washington

Myers, N., Mittermeier, R.A., Mittermeier, C.G., da Fonseca, G.A.B. & Kent, J. 2000. Biodiversity hotspots for conservation priorities. Nature 403, 853-858Powell M., A. Mills, and C. Marais. Undated. Carbon sequestration and restoration: challenges and opportunities in subtropical thicket. Unpublished document available at: <a href="https://www2.dawf.za">www2.dawf.za</a>.

Roux, D., de Moor, F. Cambray, J. & Barber-James, H. 2002. Use of landscape-level river signatures in conservation planning: a South African case study. Conservation Ecology 6, 6-15.SANBI. 2009. Indicator Report: Terrestrial Habitat Loss 2009/2/4. Unpublished report.

SRK Consulting. 2009. Final Conservation Assessment and Plan for the Nelson Mandela Bay Municipality. SRK Consulting Report Number 367380/5, SRK Consulting, Port Elizabeth.

Smith, N. & Wilson, S.L. 2002. Changing land use trends in the thicket biome: pastoralism to game farming. Terrestrial Ecology Research Unit Report 38, University of Port Elizabeth, Port Elizabeth.

SSI 2009. An invasive alien plant clearing strategy for the Nelson Mandela Bay Municipality. SSI Engineers & Environmental Consultants Report, Johannesburg.

Stewart, W.I. 2008. A Biodiversity Conservation Assessment and Plan for the Nelson Mandela Bay Municipality, Cape Floristic Region, South Africa. M.Sc. Dissertation, Nelson Mandela Metropolitan University, Port Elizabeth.

UNESCO 2009. Alexandria Coastal Dunefields. World Heritage Programme website: <a href="http://whc.unesco.org/en/tentativelists/5453/">http://whc.unesco.org/en/tentativelists/5453/</a>

Page 58

# **Appendices**

# Appendix A: Systematic Conservation Assessment and Plan for the Nelson Mandela Bay Municipality (SRK Consulting, 2010)

STEWreeb

Page 59

# **Appendix B: Extensive Biodiversity Profile**

Page 60

# Appendix C: A3 map of Critical Biodiversity Areas

STEWreeb

Draft Bioregional Plan 2 Dec 10 for gazetting (2014-final)

Page 61

# **SRK Report Distribution Record**

Report No.	400919/2
Copy No.	Electronic

Name/Title	Company	Сору	Date	Authorised by
Port Elizabeth Main Library	NMBM	1	14 April 2014	R. Gardiner
Uitenhage Town Library	NMBM	2	14 April 2014	R. Gardiner
Despatch Library	NMBM	3	14 April 2014	R. Gardiner
NMBM website	NMBM	Electronic	14 April 2014	R. Gardiner
Government Gazette	Government of the Republic of South Africa	Electronic	To be confirmed	R. Gardiner

Approval Signature:



This report is protected by copyright vested in SRK (SA) (Pty) Ltd. It may not be reproduced or transmitted in any form or by any means whatsoever to any person without the written permission of the copyright holder, SRK.

#### NOTICE - CHANGE OF TELEPHONE NUMBERS: GOVERNMENT PRINTING WORKS

As the mandated government security printer, providing world class security products and services, Government Printing Works has adopted some of the highly innovative technologies to best serve its customers and stakeholders. In line with this task, Government Printing Works has implemented a new telephony system to ensure most effective communication and accessibility. As a result of this development, our telephone numbers will change with effect from 3 February 2014, starting with the Pretoria offices.

The new numbers are as follows:

Switchboard : 012 748 6001/6002

Advertising : 012 748 6205/6206/6207/6208/6209/6210/6211/6212

Publications Enquiries: 012 748 6052/6053/6058 GeneralEnquiries@gpw.gov.za

Maps : 012 748 6061/6065 <u>BookShop@gpw.gov.za</u>

Debtors : 012 748 6060/6056/6064 PublicationsDebtors@gpw.gov.za

Subscription: 012 748 6054/6055/6057 Subscriptions@gpw.gov.za

• SCM : 012 748 6380/6373/6218

Debtors : 012 748 6236/6242

Creditors
 012 748 6246/6274

Please consult our website at www.gpwonline.co.za for more contact details.

The numbers for our provincial offices in Polokwane, East London and Mmabatho will not change at this stage.

Printed by and obtainable from the Government Printer, Bosman Street, Private Bag X85, Pretoria, 0001.

Tel: (012) 748 6052, 748 6053, 748 6058

Also available at the Legal Advisory Services, **Province of the Eastern Cape**, Private Bag X0047, Bisho, 5605. Tel. (040) 635-0052